

Accepted Manuscript

Accepted Manuscript (Uncorrected Proof)

Title: Intra-Accumbal Administration of Mecamylamine Reverses the Effects of Cytisine on the Operant Oral Self-Administration of Ethanol in Rats

Running Title: Mecamylamine Reduced EtOH Self-Administration

Authors: Juan C Jimenez¹, Rosa I Ruiz Garcia¹, Benita Cedillo-Ildefonso¹, David Hernandez¹, Florencio Miranda^{1,*}

1. *Facultad de Estudios Superiores Iztacala Av. De los Barrios 1 Los Reyes Iztacala Tlalnepantla Estado de México, México.*

***Corresponding Author:** Florencio Miranda, Facultad de Estudios Superiores Iztacala Av. De los Barrios 1 Los Reyes Iztacala Tlalnepantla Estado de México, México. Email: fmirandah@yahoo.com

To appear in: **Basic and Clinical Neuroscience**

Received date: 2022/10/12

Revised date: 2023/02/14

Accepted date: 2023/02/12

This is a “Just Accepted” manuscript, which has been examined by the peer-review process and has been accepted for publication. A “Just Accepted” manuscript is published online shortly after its acceptance, which is prior to technical editing and formatting and author proofing. *Basic and Clinical Neuroscience* provides “Just Accepted” as an optional and free service which allows authors to make their results available to the research community as soon as possible after acceptance. After a manuscript has been technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as a published article. Please note that technical editing may introduce minor changes to the manuscript text and/or graphics which may affect the content, and all legal disclaimers that apply to the journal pertain.

Please cite this article as:

Jimenez, J.C., Ruiz Garcia, R.I., Cedillo-Ildefonso, B., Hernandez, D., Miranda, F. Intra-Accumbal Administration of Mecamylamine Reverses the Effects of Cytisine on the Operant Oral Self-Administration of Ethanol in Rats. *Basic and Clinical Neuroscience*. Just Accepted publication Jul. 10, 2023. Doi: <http://dx.doi.org/10.32598/bcn.2023.4730.1>

DOI: <http://dx.doi.org/10.32598/bcn.2023.4730.1>

ABSTRACT

Introduction: It has been suggested that nicotinic acetylcholine receptors (nAChRs) expressed in the ventral tegmental area and the nucleus accumbens (nAcc) play a modulatory role in the effects of drugs of abuse. This research was designed to assess the effects of intra-accumbal administration of the nAChR antagonist mecamylamine and agonist cytosine on the operant oral self-administration of ethanol (EtOH) in rats.

Methods: Male Wistar rats were water deprived for 24 h and then trained to lever-press for EtOH reinforcement on a FR1 schedule for three sessions; thereafter, the number of responses in the FR schedule was increased to 3 until the response rate remained stable at 80%. After this training, the rats received an intra-accumbal injection of the nAChR antagonist mecamylamine (0.0, 1.25, 2.5, 5.0 µg), then nAChR agonist cytosine (0.0, 0.8, 1.6, 3.2 µg) or the combination of mecamylamine (0.0, 1.25, 2.5, 5.0 µg) and cytosine (3.2 µg) before being provided access to EtOH on a FR3 schedule.

Results: The data showed that intra-accumbal administration of mecamylamine reduced operant oral self-administration of EtOH, whereas cytosine increased operant oral self-administration of EtOH. This effect was reversed by mecamylamine.

Conclusions: These findings suggest that nAChRs in the nAcc may modulate the operant oral self-administration of EtOH in rats.

Keywords: Nicotinic acetylcholine receptors, Ethanol, Nucleus accumbens, Ethanol self-administration.

Highlights

The mesocorticolimbic dopamine (DA) system plays a role in the addiction to EtOH
DA release into the nAcc by VTA presynaptic DA terminals is controlled by nicotinic acetylcholine receptors (nAChRs)

Activation or blockade of nAChRs in the nAcc could also have a modulatory effect on EtOH-related behaviors

Intra-nAcc administration of nAChR ligands modulates oral EtOH self-administration

Plain language

Addiction to ethyl alcohol or ethanol (EtOH), also known as alcoholism, is a major public health problem around the world. The dopamine (DA) pathway in the brain reward system plays a key role in mediating the addictive effects of EtOH and other drugs of abuse. It has been suggested that DA release into the nucleus accumbens (nAcc) by ventral tegmental area presynaptic DA terminals is controlled by nicotinic acetylcholine receptors (nAChRs). The current study examined the effects of intra-accumbal administration of the nAChR antagonist mecamylamine and agonist cytosine on the operant oral self-administration of ethanol (EtOH) in rats. We found that mecamylamine reduced oral self-administration of EtOH, whereas cytosine showed an increase. The effects of cytosine were reversed by mecamylamine.

1.-INTRODUCTION

Ethyl alcohol or ethanol (EtOH) abuse is a major public health problem around the world (World Health Organization, 2018), including in Mexico (ENCODAT, 2017). Therefore, it is of fundamental importance to study the neurobiology of EtOH-related behaviors. The mesocorticolimbic dopamine (DA) system plays a key role in mediating the addictive effects of EtOH and other drugs of abuse. This system comprises DAergic neurons in the ventral tegmental area (VTA) that project their axons to the nucleus accumbens (nAcc), prefrontal cortex, amygdala and other limbic structures. EtOH interacts with several neurotransmission systems to produce its reinforcing effects, such as the glutamate (Woodward, 2000), serotonin (Sari, 2013; Yan, Zheng, Feng, & Yan, 2005), norepinephrine (Weinshenker, Rust, Miller, & Palmiter, 2000), glycine (Söderpalm, Lidö, & Ericson, 2017) and GABA (Koob, 2004) systems.

DA release into the nAcc by VTA presynaptic DA terminals is controlled by several other neurotransmitters, such as acetylcholine (ACh). ACh is released from cholinergic interneurons (iACh) within the nAcc and acts on nicotinic ACh receptors (nAChRs) and muscarinic ACh receptors. nAChRs are expressed on DAergic nerve endings. Optogenetic activation of iACh elicits DA release in the nAcc (Cachope, Mateo, Mathur, Irving, & Wang, 2012) and could modulate the effects of drugs of abuse. For example, it has been reported that the nAChR antagonist mecamylamine reduces cocaine self-administration in rats (Hansen & Mark, 2007) and blocks the acquisition of reinforcement for remifentanyl, a potent synthetic opioid (Crespo, Sturm, Saria, & Zernig, 2006), nicotine-induced locomotor activity (Bevins & Besheer, 2001) and nicotine self-administration (Ding, Gao, Sentir, & Tan, 2021). In addition, it has also been reported that the nAChR partial agonist varenicline decreases methamphetamine intake in rats (Kangiser, Dwoskin, Zheng, Crooks, & Stairs, 2018).

These data suggest that the activation or blockade of nAChRs in the nAcc could also have a modulatory effect on EtOH-related behaviors. Therefore, this study was designed to evaluate the effects of cytosine and mecamylamine, a nAChR agonist and antagonist, respectively, on operant oral EtOH self-administration in rats.

2.- MATERIALS AND METHODS

Subjects

Male Wistar rats weighing 250-300 g at the beginning of the experiment were used. The rats were individually housed in standard plastic rodent cages in a colony room maintained at a temperature of 21°C (\pm 1°C) under a 12 h light/dark cycle (lights on at 6:00 am) and had continuous access to water and food (Teklad LM485 Rat Diet from Harlan, Mexico City, Mexico). All experiments were conducted during the light phase (between 11:00 am and 1:00 pm). Animal care and handling procedures were conducted in accordance with the Official Mexican Norm (NOM-062-ZOO-1999) entitled "Technical Specifications for the Production, Care, and Use of Laboratory Animals".

Apparatus

Training and testing were conducted in five modular operant test chambers (30.5 cm long by 24 cm wide by 25 cm high) housed inside sound-attenuating cabinets with a ventilation fan (MED-008-D1 model; Med Associates, St. Albans, VT). Each chamber was equipped with two 4.5 cm wide retractable levers elevated 6.5 cm above the floor of the chamber and a stimulus light located 6 cm above each lever. A force of 0.25 N was required to activate the microswitch, and responses were reinforced with access to water (0.01 ml) or EtOH (0.01 ml). The availability of liquids was signaled by a light on the wall in front of the drinking cup that would light up for the duration of liquid delivery. The chambers were connected to a PC equipped with software for programming sessions as well as data recording.

Drugs

Ethyl alcohol or ethanol (99.91%) was purchased from J. T. Baker (Mexico City, Mexico), and cytosine ((1*R*,5*S*)-1,2,3,4,5,6-hexahydro-1,5-methano-8*H*-pyrido[1,2-*a*][1,5]diazocin-8-one) and mecamlamine ((1*R*,2*S*,4*S*)-*rel*-*N*,2,3,3-tetramethylbicyclo [2.2.1]heptan-2-amine hydrochloride) were purchased from Tocris Bioscience (Ballwin, MO, USA). Ethanol was diluted in filtered tap water (12% v/v). Cytosine and mecamlamine were dissolved in saline. All drugs were prepared fresh daily.

Surgical procedure for intracerebral cannulation

Rats were anesthetized with a ketamine:xylazine mixture (22.5 mg/kg:112.5 mg/kg, ip) and placed in a stereotaxic frame (Stoelting, Wood Dale, Illinois, USA). A guide cannula (23 G, 22 mm long; BD Precision Glide, Becton Dickinson and Co., Mexico) was unilaterally implanted in the right hemisphere of the rat brain (unilateral implantation produces less damage to brain tissue than bilateral implantation. In addition, it has been reported that in some behavioral paradigms did not find evidence for a superiority of one or other hemisphere [Schildein, Huston, & Schwarting, 2002]) 0.2 mm above the nAcc shell (AP +1.4 mm, ML 0.6 mm, DV 4.6 mm relative to bregma, 22° angle from vertical) according to the stereotaxic atlas of Paxinos and Watson (2007). After surgery, all animals were injected with an antibiotic (benzathine penicillin 300,000 UI/kg im) to prevent infection. The rats were allowed a 5-days postoperative recovery period.

Histology

After intra-nAcc administration of the cholinergic ligands, the rats were killed with a lethal dose of halothane (5%), and the brains were removed from the skull and preserved in 10% formaldehyde/saline solution for 1 week. The brains were then frozen and cut into 300 µm serial coronal slices on a vibratome. The injection sites were verified under a light microscope.

Behavioral procedures

Access to one bottle of water or EtOH solution

Rats were trained for 7 days to drink water in a 30 min period. After this training, the rats were placed in the experimental cages, where they had access to one bottle of water or EtOH solution for 30 min.

Training and testing procedure for operant oral EtOH self-administration

The timeline of the general procedure is shown in Fig. 1. The training and testing procedures were similar to those previously described (Jimenez et al., 2022). This experimental procedure leads to stable baseline of responses under FR schedule of reinforcement of oral self-administration of EtOH. For the first 7 days, tap water availability was restricted to 30 min of access to one bottle of water in the home-cages. For the next 4 days, the rats had 30 min access to one bottle of 12% EtOH solution. After this training, the rats were water deprived for 24 h and then trained in operant conditioning chambers to press a lever for water reinforcement on a fixed-ratio 1 (FR1) schedule in 30-min daily sessions for 3 consecutive days. After these training sessions, the rats were provided access to 12% EtOH solution on a FR1 schedule of reinforcement for 3 consecutive days. Thereafter, the number of responses in the FR schedule was increased to 3, i.e., each third response was reinforced with 12% EtOH (0.01 ml), until the response rate remained stable at 80% for 3 consecutive days. Once a stable response ratio was established, the effects of cholinergic receptor ligands were evaluated by injecting the rats with dose 1 of drug 1 and providing them access to 12% EtOH solution on a FR3 schedule of reinforcement for a 30-min session. After this test, reinstatement sessions were performed under conditions identical to those experienced by the rats previous to the test session. A cycle of 12% EtOH solution reinforcement on a FR3 schedule-test-12% EtOH solution reinforcement on a FR3 schedule was applied until all doses of the drugs were evaluated.

Acute effects of intra-nAcc administration of cholinergic receptor ligands on water intake

To investigate the acute effects of cytosine and mecamlamine on water intake, we conducted an initial experiment. The effects of different doses of cytosine (0.0, 0.8, 1.6, 3.2 μ g) and mecamlamine (0.0, 1.25, 2.5, 5.0 μ g) were assessed in different groups of rats ($n=6$) according to the procedure described in the access to one bottle of water section. The doses of cytosine and mecamlamine were chosen according on previous behavioral studies in which these drugs had been injected either into the nAcc or the VTA (cytosine: Reavill & Stolerman, 1990; mecamlamine: Collins, Aitken, Greenfield, Ostlund, & Wassum, 2016; Pratt & Kelley, 2004; Schiltein, Huston, & Schwarting, 2002).

Effects of intra-nAcc administration of a cholinergic receptor agonist and antagonist on operant oral EtOH self-administration

In this experiment, a within-subjects design in which a group of rats ($n=10$) was trained according to the procedure described in the training and testing procedure for operant oral EtOH self-administration section was used. After the response rate for 12% EtOH remained stable at 80% on a FR3 schedule for 3 consecutive days, the rats underwent implantation of a guide cannula according to the surgical procedure for intracerebral cannulation described above. After the surgery, the rats were allowed a 5-days recovery period. Following recovery, the rats were retrained to lever press for 12% EtOH solution reinforcement on a FR3 schedule until the response rate remained stable at 80% for 3 consecutive days. The effects of intra-nAcc administration of different doses of cytosine (0.0, 0.8, 1.6, 3.2 μg), mecamlamine (0.0, 1.25, 2.5, 5.0 μg) and coadministration of cytosine (3.2 μg) and mecamlamine (0.0, 1.25, 2.5, 5.0 μg) were tested, with one dose being evaluated per test session. A pharmacological strategy to confirm if the cytosine is acting through nAChRs is to administer an antagonist such as mecamlamine that binds to nAChRs. The drugs were administered by microinjection with calibrated polyethylene tubing connected to Hamilton syringes, and the animals were placed in the chambers for data recording 10 min later. The rate of infusion was 0.5 $\mu\text{l}/\text{min}$. After administered of each tested dose, the rats were retrained to lever press for 12% EtOH solution reinforcement on a FR3 schedule until the response rate remained stable at 80% for 3 consecutive days. The dose to be tested was randomly chosen.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics 22 software. The data were expressed as the mean (\pm standard error) of water or EtOH intake (ml) or mean number of lever presses (\pm standard error). The data obtained during the evaluation of the acute effects of cytosine and mecamlamine on water intake were analyzed using one-way analysis of variance (ANOVA). The mean number of lever pressed in the training and testing procedure for operant oral EtOH self-administration was analyzed using one-way ANOVA for repeated measures. When the ANOVA results were significant, Bonferroni's test ($p<0.05$) was used to perform post hoc comparisons.

3.- RESULTS

Acute effects of intra-accumbal administration of cholinergic receptor ligands on water intake

The results of this experiment revealed that neither cytosine ($F[4,29]=0.850$, $p=0.507$) nor mecamlamine ($F[4,29]=0.486$, $p=0.746$) altered water intake (data not shown).

Effects of intra-nAcc administration of cholinergic receptor agonist and antagonist on operant oral EtOH self-administration behavior

As shown in Table 1A-C, training sessions on FR schedules of reinforcement produced stable responses; the response rate for water was stable on a FR1 schedule ($F[2,18]=2.180$, $p=0.142$), the response rate for 12% EtOH was stable on a FR1 schedule ($F[2,18]=1.291$, $p=0.299$), and the response rate for 12% ethanol was stable on a FR3 schedule ($F[2,18]=0.639$, $p=0.539$). The

response rate to 12% EtOH on a FR3 schedule after the 5-days postsurgical recovery period is shown in Table 1D. As noted, the FR3 schedule of reinforcement produced a stable response rate ($F[2,16]=0.129$, $p=0.880$). Intra-nAcc administration of the nAChR antagonist mecamylamine decreased the operant oral self-administration of 12% EtOH (see Fig. 2A). One-way ANOVA for repeated measures revealed significant differences ($F[3,21]=6.167$, $p=0.004$). The Bonferroni test revealed that the effect of 5.0 μg of mecamylamine was different from the effect of 0.0 and 1.25 μg of mecamylamine. Intra-accumbal administration of the nAChR agonist cytosine increased the operant oral self-administration of EtOH (see Fig. 2B). One-way ANOVA for repeated measures revealed significant differences ($F[3,21]=11.064$, $p=0.001$). The Bonferroni test revealed that the effect of 3.2 μg of cytosine was different from that of the other doses of cytosine. Administration of a fixed dose of nAChR agonist cytosine (3.2 μg) in combination with various doses of the nAChR antagonist mecamylamine produced a dose-dependent decrease in the operant oral self-administration of 12% EtOH ($F[3,21]=7.606$, $p=0.001$). The Bonferroni test revealed that the effect of 5.0 μg of mecamylamine combined with cytosine (3.2 μg) was different from the effect of 0.0 and 1.25 μg of mecamylamine and that the effect of 2.5 μg of mecamylamine combined with 3.2 μg of cytosine was different from the effect of 0.0 of mecamylamine (see Fig. 2C). Figure 2D shows the histological localization of the injection sites in the shell of the nAcc. Rats in which the cannula was implanted outside the nAcc were excluded from the analyses (1/9).

4.- DISCUSSION

The purpose of the present study was to examine the effects of intra-nAcc administration of a nAChR antagonist and agonist, i.e., mecamylamine and cytosine, respectively, on operant oral self-administration of EtOH in rats. We found that rats learned to lever-press for 12% EtOH reinforcement and showed a stable operant response rate before and after cannula implantation into the nAcc shell (see table 1) without the use of initiation procedures for EtOH consumption, such as sucrose fading. In the present study, the operant oral EtOH self-administration was modeled as described in other studies (Blegen et al., 2018; Carnicella, Yowell, & Ron, 2011; Jimenez et al., 2022; Peana et al., 2014; Simms, Bito-Onon, Chatterjee, & Bartlett, 2010; Viudez-Martínez et al., 2018). This animal model of voluntary oral EtOH self-administration has been shown to be a useful tool to study the behavioral, neurochemical and cellular mechanisms underlying the addictive properties of EtOH and other drugs of abuse (Blegen et al., 2018; de Siqueira Umpierrez et al., 2022; Fernandes et al., 2020; Haile, Carper, Nolen, & Kosten, 2021). Before beginning the main experiments, the acute effects of intra-nAcc administration of mecamylamine and cytosine on water intake were studied in separated groups of rats in an initial experiment. We observed that neither mecamylamine nor cytosine altered drinking behavior at the doses used in this research.

In the current research, we also found that intra-nAcc administration of the nAChR antagonist mecamylamine reduced operant oral EtOH self-administration, while the nAChR agonist cytosine increased operant oral EtOH self-administration. This effect was reversed by the administration of mecamylamine. These observations suggest that nAChRs in the nAcc may be involved in the modulation of operant oral EtOH self-administration. Antagonists of specific

receptor subtypes are often used to confirm the mechanism associated with agonist-induced changes in some behaviors. In the present study, the intra-nAcc administration of mecamylamine, a noncompetitive nAChR antagonist, reduced the effects of cytosine on operant oral EtOH self-administration in rats. Although mecamylamine and cytosine are not selective for nAChRs, they are common pharmacological ligands employed in behavioral and neurobiological studies (Gotti & Clementi, 2021; Hendrickson, Zhao-Shea, & Tapper, 2009).

The above behavioral results are consistent with previous reports demonstrating that nAChR ligands modulate some EtOH-induced behaviors. For instance, systemic administration of mecamylamine (0-8 mg/kg) reduced EtOH self-administration following a sucrose fading procedure in C57BL/6J mice (Ford et al., 2009). Similarly, systemic administration of mecamylamine (1.25, 2.5, and 5.0 mg/kg) reduced operant oral EtOH self-administration and blocked the deprivation-induced increase in alcohol consumption (Kuzmin, Jerlhag, Liljequist, & Engel, 2009). It has also been reported that intra-accumbal administration of mecamylamine reduces EtOH self-administration but not sucrose self-administration (Nadal, Chappell, & Samson, 1998). Systemic administration of mecamylamine (1.0 and 2.0 mg/kg) also significantly reduced EtOH consumption in a limited access procedure (Lê, Corrigall, Harding, Juzytsch, & Li, 2000). Another study reported that mecamylamine reduced EtOH consumption and EtOH preference in a two-bottle choice test procedure after mecamylamine was administered both intermittently and daily (Farook, Lewis, Gaddis, Littleton, & Barron, 2009).

In the current research, intra-nAcc administration of the nAChR agonist cytosine, a partial agonist, at a dose of 3.2 μ g increased operant oral EtOH self-administration. In contrast, it has been reported that intraperitoneal administration of cytosine at a dose of 3.0 mg/kg reduces EtOH consumption in a drinking-in-the-dark procedure in mice (Hendrickson et al., 2009). Other studies have also reported that cytosine decreases EtOH-related behaviors. For example, intraperitoneal cytosine administration (1.5 mg/kg) reduced EtOH in a preference test after 24 h of concurrent access to 15% and 30% EtOH (Bell, Eiler, Cook, & Rahman, 2009). It has also been reported that intraperitoneal administration of cytosine (0.5 and 1.0 mg/kg) or varenicline (0.5 and 1.0 mg/kg) for three consecutive days reduces EtOH preference after continuous access to EtOH for four weeks in rats (Sotomayor-Zarate et al., 2013). Varenicline, a partial agonist of $\alpha 4\beta 2$ nAChR, is a synthetic derivative of cytosine (Canu Boido & Sparatore, 1999) that has been shown to reduce EtOH intake. For example, acute subcutaneous administration of varenicline (1.0 or 2.0 mg/kg) dose-dependently attenuated oral EtOH self-administration and EtOH consumption in a two-bottle choice procedure in rats (Steensland, Simms, Holgate, Richards, & Bartlett, 2007). Additionally, it has been reported that intraperitoneal administration of varenicline (0.5, 1.0, 2.0 mg/kg) dose-dependently reduces EtOH consumption in rats under a drinking-in-the-dark procedure, but it also reduces saccharin intake (Kamens, Silva, Peck, & Miller, 2018). In contrast with these studies, another study showed that systemic administration of varenicline at lower doses (0.56 and 1.0 mg/kg) increased EtOH self-administration following a sucrose fading procedure in Lewis rats without affecting responses to food. However, higher doses of varenicline decreased the response to EtOH and food (Ginsburg & Lamb, 2013).

The reason for this discrepancy between our study of intra-nAcc administration of cytosine and some of the above-cited studies is not clear. One possible hypothesis is differences in the route of drug administration. As noted above, systemic administration of nAChR agonists such as cytosine and varenicline reduced EtOH-related behaviors, whereas cytosine increased operant oral EtOH self-administration when directly administered into the nAcc in the present study. Systemic administration of the nAChR agonist cytosine can activate nAChRs expressed on different neurons in the brain reward system to reduce EtOH-related behaviors instead of increasing such behaviors. $\alpha 4\beta 2$ nAChRs (which are activated by Ach and nicotine and are involved in modulating the rewarding effects of EtOH) are expressed on DAergic neurons projecting to the nAcc and on local GABA interneurons in the VTA and modulate the firing patterns of DA neurons (Mameli-Engvall et al., 2006; Maurer & Schmidt, 2019). In addition, $\alpha 4\beta 2$ nAChRs are also expressed on DA terminals in the nAcc (Feduccia, Chatterjee, & Bartlett, 2012; Grady et al., 2007). Therefore, it can be speculated that systemic administration of cytosine can lead to activation of $\alpha 4\beta 2$ nAChRs expressed on all these neurons, including those expressed on the cell bodies of GABAergic interneurons in the VTA, to inhibit EtOH-related behaviors, whereas intra-nAcc administration of cytosine promotes EtOH-related behaviors, but this possibility remains to be evaluated. Another possible hypothesis accounting for the discrepancy between the above-cited data and the results of the present study is the pharmacological profile of cytosine. Cytosine is an alkaloid with partial agonist activity at $\alpha 4\beta 2$ nAChRs, and it has been shown to reduce rather than increase EtOH-related behaviors in several animal models (see above). A partial agonist does not produce the maximal response and can act as an antagonist in the presence of a full agonist depending on the dose. It has been reported that cytosine (3.0 mg/kg) partially substitutes for nicotine (59%) in rats trained to discriminate nicotine (0.6 mg/kg) from saline (Radchenko, Dravolina, & Beshpalov, 2015). In addition, it has also been reported that intraperitoneal administration of a high dose of cytosine (3.0 mg/kg) partially generalizes (23% of nicotine-appropriate lever presses) and antagonizes nicotine-induced discriminative stimulus effects in rats (LeSage, Shelley, Ross, Carroll, & Corrigan, 2009). In general, the discrepancy between the effects of intra-nAcc administration of cytosine in our study and some of the above-cited data may be related to differences in drug administration procedures, dosage regimen or procedural variables.

Inhibition or activation of $\alpha 4\beta 2$ nAChRs in the nAcc might be involved in the modulation of the operant oral self-administration of EtOH observed in the present study. This speculation is based on the following findings. First, the mesocorticolimbic DA system, particularly the projections from the VTA to the nAcc, is important for EtOH-related behaviors (Rodd et al., 2004; Weiss, Lorang, Bloom, & Koob, 1993). Second, $\alpha 4\beta 2$ nAChRs, which are expressed in DA terminals in the nAcc (Feduccia et al., 2012; Grady et al., 2007), play a modulatory role in DA release in the nAcc and, as a consequence, affect EtOH-related behaviors. DA release from VTA DA terminals in the nAcc is under the control of different receptors, such as D2 autoreceptors and $\alpha 4\beta 2$ nAChRs. Some studies have shown that stimulation of those receptors modulates DA release in the nAcc. For instance, optogenetic stimulation of nAcc iACh produced the DA release in nAcc presynaptic DA terminals that express $\alpha 4\beta 2$ nAChRs (Cachope et al., 2012). Furthermore, inhibition of iACh firing using an opioid agonist decreased the frequency of spontaneous DA transients in the nAcc (Yorgason, Zeppenfeld, & Williams,

2017). Consistent with this finding, behavioral studies involving manipulation of $\alpha 4\beta 2$ nAChRs have provided additional support for the modulatory role of $\alpha 4\beta 2$ nAChRs on EtOH-related behaviors. Nadal et al. (1998) reported that intra-accumbal administration of the $\alpha 4\beta 2$ nAChR antagonist mecamylamine reduced EtOH self-administration but not sucrose self-administration in rats. $\alpha 4\beta 2$ nAChRs are not only involved in EtOH-related behaviors but are also involved in drug abuse-related behaviors. For instance, it has been reported that pretreatment with mecamylamine (0.3, 1.0, 3.0 mg/kg) reduces cocaine and nicotine self-administration in rats (Blokhina, Kashkin, Zvartau, Danysz, & Bessalov, 2005). Another study showed that mecamylamine reduced cocaine self-administration but not food self-administration (Levin et al., 2000). Although in these studies, the $\alpha 4\beta 2$ nAChR antagonist mecamylamine was administered systemically, it was able to reach and block nAChRs expressed on different neurons in the brain reward system, including the nAcc, to reduce drug abuse-related behaviors, although more research is needed before coming to a final conclusion regarding the expression of $\alpha 4\beta 2$ nAChRs in different regions of the brain reward system and their involvement in EtOH-related behaviors.

In summary, our results showed that intra-accumbal administration of $\alpha 4\beta 2$ nAChR ligands modulated operant oral EtOH self-administration. These data provide further evidence that $\alpha 4\beta 2$ nAChRs may modulate drug abuse-induced behaviors, particularly the reinforcing effects of EtOH.

Author contributions

Conceptualization and supervision: FM; Methodology: RIR, BC-I and DH; Data analysis: JCJ; Written-original draft: FM and JCJ; Investigation: All authors; Funding and resources: FM; Approving the final version of manuscript: All authors

Acknowledgments

This study was supported by grant IN305420 from PAPIIT-UNAM (México). The authors declare no conflicts of interest.

5.- REFERENCES

- Bell, R. L., Eiler, B. J., 2nd, Cook, J. B., & Rahman, S. (2009). Nicotinic receptor ligands reduce ethanol intake by high alcohol-drinking HAD-2 rats. *Alcohol (Fayetteville, N.Y.)*, 43(8), 581–592. <https://doi.org/10.1016/j.alcohol.2009.09.027>
- Bevins, R. A., & Besheer, J. (2001). Individual differences in rat locomotor activity are diminished by nicotine through stimulation of central nicotinic acetylcholine receptors. *Physiology & behavior*, 72(1-2), 237–244. [https://doi.org/10.1016/s0031-9384\(00\)00413-3](https://doi.org/10.1016/s0031-9384(00)00413-3)
- Blegen, M. B., da Silva E Silva, D., Bock, R., Morisot, N., Ron, D., & Alvarez, V. A. (2018). Alcohol operant self-administration: Investigating how alcohol-seeking behaviors predict drinking in mice using two operant approaches. *Alcohol (Fayetteville, N.Y.)*, 67, 23–36. <https://doi.org/10.1016/j.alcohol.2017.08.008>
- Blokhina, E. A., Kashkin, V. A., Zvartau, E. E., Danysz, W., & Bessalov, A. Y. (2005). Effects of nicotinic and NMDA receptor channel blockers on intravenous cocaine and nicotine self-administration in mice. *European neuropsychopharmacology: the journal of the European College of Neuropsychopharmacology*, 15(2), 219–225. <https://doi.org/10.1016/j.euroneuro.2004.07.005>
- Cachope, R., Mateo, Y., Mathur, B. N., Irving, J., Wang, H. L., Morales, M., Lovinger, D. M., & Cheer, J. F. (2012). Selective activation of cholinergic interneurons enhances accumbal phasic dopamine release: setting the tone for reward processing. *Cell reports*, 2(1), 33–41. <https://doi.org/10.1016/j.celrep.2012.05.011>
- Canu Boido, C., & Sparatore, F. (1999). Synthesis and preliminary pharmacological evaluation of some cytosine derivatives. *Farmaco (Societa chimica italiana: 1989)*, 54(7), 438–451. [https://doi.org/10.1016/s0014-827x\(99\)00049-x](https://doi.org/10.1016/s0014-827x(99)00049-x)
- Carnicella, S., Yowell, Q. V., & Ron, D. (2011). Regulation of operant oral ethanol self-administration: a dose-response curve study in rats. *Alcoholism, clinical and experimental research*, 35(1), 116–125. <https://doi.org/10.1111/j.1530-0277.2010.01328.x>
- Collins, A. L., Aitken, T. J., Greenfield, V. Y., Ostlund, S. B., & Wassum, K. M. (2016). Nucleus accumbens acetylcholine receptors modulate dopamine and motivation. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*, 41(12), 2830–2838. <https://doi.org/10.1038/npp.2016.81>
- Crespo, J. A., Sturm, K., Saria, A., & Zernig, G. (2006). Activation of muscarinic and nicotinic acetylcholine receptors in the nucleus accumbens core is necessary for the acquisition of

- drug reinforcement. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 26(22), 6004–6010. <https://doi.org/10.1523/JNEUROSCI.4494-05.2006>
- de Siqueira Umpierrez, L., Freese, L., Almeida, F. B., Costa, P. A., Fernandes, P. R., Nin, M. S., de Souza, M. F., & Barros, H. (2022). Effects of neonatal dopaminergic lesion on oral cocaine self-administration in rats: Higher female vulnerability to cocaine consumption. *Pharmacology, biochemistry, and behavior*, 212, 173315. <https://doi.org/10.1016/j.pbb.2021.173315>
- Ding, Z. M., Gao, Y., Sentir, A. M., & Tan, X. (2021). Self-Administration of Cotinine in Wistar Rats: Comparisons to Nicotine. *The Journal of pharmacology and experimental therapeutics*, 376(3), 338–347. <https://doi.org/10.1124/jpet.120.000367>
- ENCODAT (2017). *Encuesta Nacional de Consumo de Drogas, Alcohol y Tabaco 2016-2017*, Reporte de Alcohol. Secretaría de Salud. México.
- Farook, J. M., Lewis, B., Gaddis, J. G., Littleton, J. M., & Barron, S. (2009). Effects of mecamylamine on alcohol consumption and preference in male C57BL/6J mice. *Pharmacology*, 83(6), 379–384. <https://doi.org/10.1159/000219488>
- Feduccia, A. A., Chatterjee, S., & Bartlett, S. E. (2012). Neuronal nicotinic acetylcholine receptors: neuroplastic changes underlying alcohol and nicotine addictions. *Frontiers in molecular neuroscience*, 5, 83. <https://doi.org/10.3389/fnmol.2012.00083>
- Fernandes, P. R., Almeida, F. B., da Cunha, M., Feddern, C. F., Freese, L., & Barros, H. (2020). The effects of caffeine on alcohol oral self-administration behavior in rats. *Physiology & behavior*, 223, 112966. <https://doi.org/10.1016/j.physbeh.2020.112966>
- Ford, M. M., Fretwell, A. M., Nickel, J. D., Mark, G. P., Strong, M. N., Yoneyama, N., & Finn, D. A. (2009). The influence of mecamylamine on ethanol and sucrose self-administration. *Neuropharmacology*, 57(3), 250–258. <https://doi.org/10.1016/j.neuropharm.2009.05.012>
- Ginsburg, B. C., & Lamb, R. J. (2013). Effects of varenicline on ethanol- and food-maintained responding in a concurrent access procedure. *Alcoholism, clinical and experimental research*, 37(7), 1228–1233. <https://doi.org/10.1111/acer.12085>
- Gotti, C., & Clementi, F. (2021). Cytisine and cytosine derivatives. More than smoking cessation aids. *Pharmacological research*, 170, 105700. <https://doi.org/10.1016/j.phrs.2021.105700>
- Grady, S. R., Salminen, O., Laverty, D. C., Whiteaker, P., McIntosh, J. M., Collins, A. C., & Marks, M. J. (2007). The subtypes of nicotinic acetylcholine receptors on dopaminergic

- terminals of mouse striatum. *Biochemical pharmacology*, 74(8), 1235–1246. <https://doi.org/10.1016/j.bcp.2007.07.032>
- Haile, C. N., Carper, B. A., Nolen, T. L., & Kosten, T. A. (2021). The GABA_B receptor positive allosteric modulator ASP8062 reduces operant alcohol self-administration in male and female Sprague Dawley rats. *Psychopharmacology*, 238(9), 2587–2600. <https://doi.org/10.1007/s00213-021-05881-0>
- Hansen, S. T., & Mark, G. P. (2007). The nicotinic acetylcholine receptor antagonist mecamylamine prevents escalation of cocaine self-administration in rats with extended daily access. *Psychopharmacology*, 194(1), 53–61. <https://doi.org/10.1007/s00213-007-0822-z>
- Hendrickson, L. M., Zhao-Shea, R., & Tapper, A. R. (2009). Modulation of ethanol drinking-in-the-dark by mecamylamine and nicotinic acetylcholine receptor agonists in C57BL/6J mice. *Psychopharmacology*, 204(4), 563–572. <https://doi.org/10.1007/s00213-009-1488-5>
- Jimenez, J. C., Cortés-Salazar, F., Cedillo-Ildefonso, B., & Miranda, F. (2022). La administración sistémica e intra-accumbens del agonista 5-HT_{1B} CP94253 modula la auto-administración oral de etanol en ratas. *Revista Argentina de Ciencias del Comportamiento*, 14(1), 68-81.
- Kamens, H. M., Silva, C., Peck, C., & Miller, C. N. (2018). Varenicline modulates ethanol and saccharin consumption in adolescent male and female C57BL/6J mice. *Brain research bulletin*, 138, 20–25. <https://doi.org/10.1016/j.brainresbull.2017.07.020>
- Kangiser, M. M., Dwoskin, L. P., Zheng, G., Crooks, P. A., & Stairs, D. J. (2018). Varenicline and GZ-793A differentially decrease methamphetamine self-administration under a multiple schedule of reinforcement in rats. *Behavioural pharmacology*, 29(1), 87–97. <https://doi.org/10.1097/FBP.0000000000000340>
- Koob G. F. (2004). A role for GABA mechanisms in the motivational effects of alcohol. *Biochemical pharmacology*, 68(8), 1515–1525. <https://doi.org/10.1016/j.bcp.2004.07.031>
- Kuzmin, A., Jerlhag, E., Liljequist, S., & Engel, J. (2009). Effects of subunit selective nACh receptors on operant ethanol self-administration and relapse-like ethanol-drinking behavior. *Psychopharmacology*, 203(1), 99–108. <https://doi.org/10.1007/s00213-008-1375-5>
- Lê, A. D., Corrigall, W. A., Harding, J. W., Juzysch, W., & Li, T. K. (2000). Involvement of nicotinic receptors in alcohol self-administration. *Alcoholism, clinical and experimental research*, 24(2), 155–163. <https://doi.org/10.1111/j.1530-0277.2000.tb04585.x>

- LeSage, M. G., Shelley, D., Ross, J. T., Carroll, F. I., & Corrigall, W. A. (2009). Effects of the nicotinic receptor partial agonists varenicline and cytisine on the discriminative stimulus effects of nicotine in rats. *Pharmacology, biochemistry, and behavior*, 91(3), 461–467. <https://doi.org/10.1016/j.pbb.2008.08.024>
- Levin, E. D., Mead, T., Rezvani, A. H., Rose, J. E., Gallivan, C., & Gross, R. (2000). The nicotinic antagonist mecamylamine preferentially inhibits cocaine vs. food self-administration in rats. *Physiology & behavior*, 71(5), 565–570. [https://doi.org/10.1016/s0031-9384\(00\)00382-6](https://doi.org/10.1016/s0031-9384(00)00382-6)
- Mameli-Engvall, M., Evrard, A., Pons, S., Maskos, U., Svensson, T. H., Changeux, J. P., & Faure, P. (2006). Hierarchical control of dopamine neuron-firing patterns by nicotinic receptors. *Neuron*, 50(6), 911–921. <https://doi.org/10.1016/j.neuron.2006.05.007>
- Maurer, J. J., & Schmidt, H. D. (2019). Nicotine Addiction and Alpha4beta2* Nicotinic Acetylcholine Receptors. Editor(s): Victor R. Preedy. In *Neuroscience of Nicotine*, 243–250. Academic Press.
- Nadal, R., Chappell, A. M., & Samson, H. H. (1998). Effects of nicotine and mecamylamine microinjections into the nucleus accumbens on ethanol and sucrose self-administration. *Alcoholism, clinical and experimental research*, 22(6), 1190–1198.
- Paxinos, G., & Watson, C. (2007). *The rat brain in stereotaxic coordinates*. 6th Ed. Cambridge: Academic Press.
- Peana, A. T., Muggirani, G., Spina, L., Rosas, M., Kasture, S. B., Cotti, E., & Acquas, E. (2014). Effects of *Withania somnifera* on oral ethanol self-administration in rats. *Behavioural pharmacology*, 25(7), 618–628. <https://doi.org/10.1097/FBP.0000000000000078>
- Pratt, W. E., & Kelley, A. E. (2004). Nucleus accumbens acetylcholine regulates appetitive learning and motivation for food via activation of muscarinic receptors. *Behavioral neuroscience*, 118(4), 730–739. <https://doi.org/10.1037/0735-7044.118.4.730>
- Radchenko, E. V., Dravolina, O. A., & Beshpalov, A. Y. (2015). Agonist and antagonist effects of cytisine in vivo. *Neuropharmacology*, 95, 206–14. <https://doi.org/10.1016/j.neuropharm.2015.03.019>
- Reavill, C., & Stolerman, I. P. (1990). Locomotor activity in rats after administration of nicotinic agonists intracerebrally. *British journal of pharmacology*, 99(2), 273–278. <https://doi.org/10.1111/j.1476-5381.1990.tb14693.x>

- Rodd, Z. A., Melendez, R. I., Bell, R. L., Kuc, K. A., Zhang, Y., Murphy, J. M., & McBride, W. J. (2004). Intracranial self-administration of ethanol within the ventral tegmental area of male Wistar rats: evidence for involvement of dopamine neurons. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 24(5), 1050–1057. <https://doi.org/10.1523/JNEUROSCI.1319-03.2004>
- Sari Y. (2013). Role of 5-hydroxytryptamine 1B (5-HT1B) receptors in the regulation of ethanol intake in rodents. *Journal of psychopharmacology (Oxford, England)*, 27(1), 3–12. <https://doi.org/10.1177/0269881112463126>
- Schilwein, S., Huston, J. P., & Schwarting, R. K. (2002). Open field habituation learning is improved by nicotine and attenuated by mecamylamine administered posttrial into the nucleus accumbens. *Neurobiology of learning and memory*, 77(3), 277–290. <https://doi.org/10.1006/nlme.2001.4017>
- Simms, J. A., Bito-Onon, J. J., Chatterjee, S., & Bartlett, S. E. (2010). Long-Evans rats acquire operant self-administration of 20% ethanol without sucrose fading. *Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology*, 35(7), 1453–1463. <https://doi.org/10.1038/npp.2010.15>
- Söderpalm, B., Lidö, H. H., & Ericson, M. (2017). The Glycine Receptor-A Functionally Important Primary Brain Target of Ethanol. *Alcoholism, clinical and experimental research*, 41(11), 1816–1830. <https://doi.org/10.1111/acer.13483>
- Sotomayor-Zárate, R., Gysling, K., Busto, U. E., Cassels, B. K., Tampier, L., & Quintanilla, M. E. (2013). Varenicline and cytosine: two nicotinic acetylcholine receptor ligands reduce ethanol intake in University of Chile bibulous rats. *Psychopharmacology*, 227(2), 287–298. <https://doi.org/10.1007/s00213-013-2974-3>
- Steensland, P., Simms, J. A., Holgate, J., Richards, J. K., & Bartlett, S. E. (2007). Varenicline, an $\alpha 4\beta 2$ nicotinic acetylcholine receptor partial agonist, selectively decreases ethanol consumption and seeking. *Proceedings of the National Academy of Sciences of the United States of America*, 104(30), 12518–12523. <https://doi.org/10.1073/pnas.0705368104>
- Viudez-Martínez, A., García-Gutiérrez, M. S., Navarrón, C. M., Morales-Calero, M. I., Navarrete, F., Torres-Suárez, A. I., & Manzanares, J. (2018). Cannabidiol reduces ethanol consumption, motivation and relapse in mice. *Addiction biology*, 23(1), 154–164. <https://doi.org/10.1111/adb.12495>
- Weinshenker, D., Rust, N. C., Miller, N. S., & Palmiter, R. D. (2000). Ethanol-associated behaviors of mice lacking norepinephrine. *The Journal of neuroscience: the official*

journal of the Society for Neuroscience, 20(9), 3157–3164.
<https://doi.org/10.1523/JNEUROSCI.20-09-03157.2000>

- Weiss, F., Lorang, M. T., Bloom, F. E., & Koob, G. F. (1993). Oral alcohol self-administration stimulates dopamine release in the rat nucleus accumbens: genetic and motivational determinants. *The Journal of pharmacology and experimental therapeutics*, 267(1), 250–258. PMID: 8229752
- Woodward J. J. (2000). Ethanol and NMDA receptor signaling. *Critical reviews in neurobiology*, 14(1), 69–89. <https://doi.org/10.1080/08913810008443548>
- World Health Organization (2018). Global Status Report on Alcohol 2018. World Health Organization, Department of Mental Health and Substance Abuse. Geneva, Switzerland.
- Yan, Q. S., Zheng, S. Z., Feng, M. J., & Yan, S. E. (2005). Involvement of 5-HT1B receptors within the ventral tegmental area in ethanol-induced increases in mesolimbic dopaminergic transmission. *Brain research*, 1060(1-2), 126–137. <https://doi.org/10.1016/j.brainres.2005.08.051>
- Yorgason, J. T., Zeppenfeld, D. M., & Williams, J. T. (2017). Cholinergic Interneurons Underlie Spontaneous Dopamine Release in Nucleus Accumbens. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 37(8), 2086–2096. <https://doi.org/10.1523/JNEUROSCI.3064-16.2017>

Table 1. Summary of the mean response rate during the operant oral EtOH self-administration training procedure.

Note. A within-subjects design was used in this study (n=10). FR: fixed ratio schedule of reinforcement. E^R: reinforcing stimulus. EtOH: ethanol. SEM: standard error of the mean. One rat died after surgery.

Fig. 1. Timeline of the operant oral EtOH self-administration training and testing procedure. Schematic illustration of the procedure used to establish the operant oral EtOH self-administration model. Left side: rats were trained to drink water or EtOH for 30 min. Right side: rats were trained in operant conditioning chambers to press a lever for water or EtOH reinforcement on a fixed-ratio (FR) schedule of reinforcement.

Fig. 2.A-C. Effects of the intra-accumbal administration of nicotinic cholinergic ligands on operant oral EtOH self-administration. The bars represent the mean numbers of lever presses, and the vertical lines represent the \pm SEMs of 8 rats. The asterisks (*) indicate significant differences (Bonferroni test, $p < 0.001$, after one-way ANOVA for repeated measures) between the mean number of lever presses observed after administration of nicotinic cholinergic receptor ligands and that observed in all other conditions. The crosses (†) indicate significant differences (Bonferroni test, $p < 0.001$, after one-way ANOVA for repeated measures) between the mean number of lever presses observed after administration of nicotinic cholinergic receptor antagonist mecamylamine and the condition 0.0. D. Photograph of one coronal section taken at the level of the cannula. E. The gray bar shows the trajectory of the cannula. F. Enlargement of the site of injection. G. Schematic representation of all cannula implanted in the nAcc shell. The black circles indicate histologically confirmed locations of the cannula in the nAcc shell in the animals that were included in the statistical analyses for all experiments (8/9).

Fig. 1




Table 1

		DAYS		
		1	2	3
A	FR1 (E^R = water)			
	Lever presses mean	319.91	366.32	367.32
	SEM	14.61	26.71	23.76
B	FR1 (E^R = EtOH)			
	Lever presses mean	241.22	287.92	309.71
	SEM	29.27	38.42	34.13
C	FR3 (E^R = EtOH)			
	Last 3 days (80% stability)			
	Lever presses mean	442.22	514.93	480.21
	SEM	43.74	47.44	59.71
D	FR3 (E^R = EtOH)			
	After Surgery (80% stability)			
	Lever presses mean	558.21	576.92	554.51
	SEM	68.84	66.71	58.29

Fig. 2

Fig 2.

