Accepted Manuscript

Accepted Manuscript (Uncorrected Proof)

Title: Serotonergic Modulation of Orientation Tuning of Neurons in Primary Visual Cortex of Anesthetized Mice

Running title: Serotonergic modulation of neurons in visual cortex

Authors: Sareh Rostami¹, Amin Asgharzadeh Alvar², Parviz Ghaderi³, Leila Dargahi¹, Mir-Shahram Safari^{4*}

1. Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
2. Student Research Committee, Department of Biomedical Engineering and Medical Physics, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
3. Laboratory of Sensory Processing, Brain Mind Institute, Faculty of Life Sciences, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland.
4. Neuroscience Research Center, Department of Neuroscience, Faculty of Medicine Shahid Beheshti University of Medical Sciences, Tehran, Iran.

***Corresponding Author:** Mir-Shahram Safari, Neuroscience Research Center, Department of Neuroscience, Faculty of Medicine Shahid Beheshti University of Medical Sciences, Tehran, Iran.

E-mail: safari@sbmu.ac.ir

To appear in: *Basic and Clinical Neuroscience*

Received date: 2021/01/16

Revised date: 2021/02/3

Accepted date: 2021/02/2

This is a “Just Accepted” manuscript, which has been examined by the peer-review process and has been accepted for publication. A “Just Accepted” manuscript is published online shortly after its acceptance, which is prior to technical editing and formatting and author proofing. *Basic and Clinical Neuroscience* provides “Just Accepted” as an optional and free service which allows authors to make their results available to the research community as soon as possible after acceptance. After a manuscript has been technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as a published article. Please note that technical editing may introduce minor changes to the manuscript text and/or graphics which may affect the content, and all legal disclaimers that apply to the journal pertain.

Please cite this article as:

Rostami, S., Asgharzadeh Alvar, A., Ghaderi, P., Dargahi, L., & Safari, MS. (In Press). Serotonergic Modulation of Orientation Tuning of Neurons in Primary Visual Cortex of Anesthetized Mice. *Basic and Clinical Neuroscience*. Just Accepted publication Jun. 27, 2021. Doi: <http://dx.doi.org/10.32598/bcn.2021.3180.1>

DOI: <http://dx.doi.org/10.32598/bcn.2021.3180.1>

Abstract

Introduction: Sensory processing is profoundly regulated by brain neuromodulatory systems. One of the main neuromodulators is serotonin which influences higher cognitive functions such as different aspects of perceptual processing. So, malfunction in the serotonergic system may lead to visual illusion in psychiatric disorders such as autism and schizophrenia. In this work, we examined the serotonergic modulation of visual responses of neurons to stimulus orientation in the primary visual cortex.

Methods: Eight-weeks old naive mice were anesthetized and craniotomy was done on the region of interest in primary visual cortex. Spontaneous and visual-evoked activities of neurons were recorded before and during the electrical stimulation of dorsal raphe nucleus using in vivo whole-cell patch-clamp recording. Square-wave grating of 12 orientations was presented. Data was analyzed and Wilcoxon signed-rank test, used in order to compare the data of two conditions that belong to the same neurons, with or without electrical stimulation.

Results: The serotonergic system changed orientation tuning of about 60 % recorded neurons by decreasing the mean firing rate in two independent visual response components: gain and baseline response. It also increased mean firing rate in a small number of neurons (about 20%). Beyond that, it left the preferred orientation and sensitivity of neurons unchanged.

Conclusion: However, serotonergic modulation showed a bi-directional effect; it seems to cause predominately divisive and subtractive decreases in the visual responses of the neurons in the primary visual cortex that can modify the balance between internal and external sensory signals and result in disorders.

Keywords: Serotonergic modulation, Orientation tuning, Primary visual cortex, In vivo patch-clamp recording, Dorsal raphe nucleus

Highlights

- The serotonergic system predominantly decreased mean firing rate of neurons in primary visual cortex.
- The serotonergic system decreased responses of visual cortical neurons by subtractive and divisive changes of orientation tuning.
- The serotonergic system leaves the spontaneous activity of visual cortical neurons unchanged.

Plain Language Summary

Serotonin is one of the most well-known neuromodulators involved in many physiological functions of the brain, such as sensory processing. It can play an essential role in producing perceptual psychotic episodes following the use of psychedelic drugs. Neural mechanisms of changes in cortical processing by the serotonergic system are not elucidated enough. In this study, we showed the electrical stimulation of the dorsal raphe nucleus as the main resource of projecting serotonergic neurons to the visual cortex, caused to decrease visual-evoked responses of neurons in primary visual cortex without changing the spontaneous activity. This effect may lead to an imbalance between the brain's intrinsic and stimulus-evoked activity and result in various kinds of psychiatric disorders such as visual hallucinogenic experiences in schizophrenia and autism. Therefore, it is crucial to understand the mechanisms by which serotonin affects the rapid and long-term activity of neocortical circuits. Such studies can be helpful in diagnosis and treatment of disorders related to neuromodulatory roles of the serotonergic system by providing new methods for rebalancing these intricate components.

1- Introduction

In the early steps of sensory processing, the brain's internal states modulate the neuronal representation of external stimuli (Harris & Thiele, 2011)(Aston-Jones & Cohen, 2005). Processing of information in sensory system is highly tuned by the neuromodulatory networks that are differently active in various states, which project to specific layers and regions of cortical and large areas of subcortical structures and exert feedforward and feedback effects on signaling in sensory system by different mechanisms (Cumming & Nienborg, 2016)(Jacob & Nienborg, 2018).

Serotonin is one of the major neuromodulators of the sensory systems. Serotonergic neurons in the midbrain medial and dorsal raphe nuclei frequently project to forebrain through the medial frontal bundles (Calizo et al., 2011)(Dahlström & Fuxe, 1964)(Hornung et al., 1990). Similar to some other neuromodulatory systems, these nuclei consist of different cell types, including GABAergic, glutamatergic, and peptidergic neurons (Calizo et al., 2011)(Waselus et al., 2006)(Carcea & Froemke, 2013). As already the well-known the serotonergic system is implicated in many basic functions such as controlling blood pressure, appetite and nutrition, sleep, body temperature, pain perception, anxiety, aggression, sexual behaviors, reward, mood and emotions, social cognition, learning and memory, motor control and sensory processing (Muller & Cunningham, 2020)(Pollak Dorocic, 2016)(Curzon, 1988)(Peroutka, 1991)(Lucki, 1992)(Tseng & Atzori, 2007). The system is also involved in a different kind of psychiatric disorders, including schizophrenia, ADHD, depression, autism, drug addiction, OCD, stress, eating disorders, aggression and anxiety disorders; as well as malfunction of this system is the basis for the effects of the hallucinogenic drugs (Lucki, 1998). Serotonin and its receptors can play an essential role in the production of perceptual psychotic episodes following the use of psychedelic drugs as well as in schizophrenia (González-Maeso et al., 2008)(Lesch & Waider, 2012). Therefore, it is crucial to understand the mechanisms by which serotonin affects neocortical circuits' rapid and long-term activity.

Serotonergic system has projected mainly to the primary sensory regions. It probably has an important effect in tuning sensory inputs from the early stages in the primary sensory areas and thalamic relay nuclei (Jacob & Nienborg, 2018). Serotonin is also implicated in the functional and structural regeneration of circuits in cortex. Serotonergic neurons affect areas of cortex that are involved in processing sensory information in audition, somatic sensation and vision. Their response properties can be changed by neuromodulators following long-term changes in

sensory input activity, specifically in early postnatal life (Qiang Gu, 2003)(Q Gu, 2002). In vivo and in vitro evaluation of serotonin function on cortical neurons has shown complicated patterns. Literature offers that the effect of serotonin in the cerebral cortex is to regulate cortical neurons excitability during sensory gating, determining the threshold for synaptic changes related to activity and increasing signal-to-noise ratio (Tseng & Atzori, 2007). Despite the diversity of serotonergic modulation researches on sensory processing, the more common effect is to reduce the gain of neuronal responses (Jacob & Nienborg, 2018).

There are serotonergic receptors, including an inotropic receptor (5-HT_{3A}) and different types of metabotropic receptors on inhibitory interneurons and pyramidal neurons in the cortex. Metabotropic receptors of serotonin can regulate potassium channel activity and intracellular calcium levels and interact directly or indirectly with glutamate metabotropic receptors, thus controlling synaptic transmission, neuronal excitability, and excitation-inhibition balance in the network (Bockaert et al., 2010)(William Moreau et al., 2010)(Ögren et al., 2008)(Yuen et al., 2005). Receptor 5-HT_{3A} is the main marker of the one kind of three types of inhibitory interneurons, indicating that serotonin's modulatory role in controlling inhibition of cortical neural networks would be specific and precise function.

The visual cortex's functional status is mainly regulated by non-visual input systems, that are neuromodulatory inputs from subcortical areas and can modify responses of visual input in order to overcome the threshold of anatomical and physiological changes related to activity (Tseng & Atzori, 2007). In the visual cortex, serotonin neurotransmission is assumed one of the non-visual inputs that might partly involve in the neurochemical basis of motivation, attention and excitation and influence them. One of the functions of serotonergic system in cortical processing of vision is to change response of neurons to visual sensory input. The stimulation of dorsal raphe nucleus electrically and in vivo direct serotonin application on the cortical neurons demonstrate the neuronal activity suppression or facilitation (Waterhouse et al., 1990)(Reader, 1978)(Krnjević & Phillis, 1963). Whereas non-visual inputs to the visual cortical areas are essential modulators of plasticity in visual cortex (Q Gu, 2002)(Qiang Gu, 2003), serotonin's involvement in cortical plasticity has also been confirmed in vivo and in vitro.

Previous studies showed serotonin can affect GABAergic interneurons directly in the visual cortical areas through its receptors. For example, 5-HT₃ receptor activity reduces the size and

pattern of excitation in ferret visual cortex (Roerig & Katz, 1997), probably due to an increase in GABAergic synaptic activity (Xiang & Prince, 2003).

Considering complexity of the serotonergic system function in neuronal networks, and because most works on the effect of serotonin in visual processing have been done using an extracellular recording, we would like to investigate the supra-threshold and subthreshold responses of neurons to stimulus orientation in the primary visual cortex of anesthetized mice.

2- Material and methods

2-1- Animal and craniotomy

Eight-weeks old naive mice weight 20-30 gr of either sexes were used. The animal was anesthetized by intraperitoneal urethane injection (1.5-2 mg/gr body weight). After cutting the hair of the head and mustache, it was placed inside the stereotaxic device, and the head was kept fixed. Part of the skin and muscles was removed in the parietal area. A craniotomy of 1.5 mm × 2 mm (AP: 2.5-4 mm and L: 2-4 mm) was done on the left hemisphere's primary visual cortex area (area 17). A metal head plate was planted on the skull by superglue and dental acrylic. After thinning the skull, part of it as well as the dura matter will be removed on the primary visual cortex to allow access to the cortex. The temperature of animal's body was sustained at 36-36.5 ° C with a rectal thermal probe (ATC-402, Unique Medical).

2-2- Visual stimulation

The visual stimulus includes a drifting grating stimulus using spatial frequency of 0.04 cycles/degree, a contrast of 100%, and a temporal frequency of 1 Hz. Square-wave gratings of 12 orientations in 30-degree (0-360 °) steps were moved on an LCD monitor (Flexscan L788, 19 inch, Eizo Nanao). Visual stimuli of 12 orientation patterns (2s before, 3s during and 2s after the presentation of each stimulus, respectively) were presented in shuffled random order for three times. The LCD monitor was located at a distance of 28 cm from the eyes of animal and included 80*50 degrees' field of vision.

2-3- The serotonergic system activation by electrical stimulation of dorsal raphe nucleus (DRN)

To stimulate the serotonergic projections of DRN to the primary visual cortex in the anesthetized animal, we used DRN electrical stimulation. We implanted a bipolar tungsten electrode in the brain using a micromanipulator and a stereotaxic atlas of the mouse brain (AP:

4.3 mm, L: 0, and DV: 3 mm). We have done the histological validation in order to confirm the exact location of electrode insert by destroying the area and then cutting to slices and observing under a microscope. The DRN was stimulated electrically at 400-700 μ A, and pulse frequency was 20 Hz (Mokler et al., 1998) three times during visual stimulation.

2-4- In vivo whole-cell patch-clamp recording

Whole-cell patch-clamp recording of excitatory and inhibitory neurons of layer I at a depth of 20-90 μ m and layer II/III at a depth of 200-300 μ m were performed blindly. The electrodes for recording were made from borosilicate glass capillary with filaments (inner diameter of 0.86 mm and an outer diameter of 1.5 mm), filled by a pre-prepared internal solution. The internal solution had the osmolarity of 280-290 mOsm and adjusted its pH with KOH to 7.2-7.4, which is equivalent to the intracellular environment, and its compounds were (in mM): CaCl₂, 0.1; MgATP, 4; K-Gluconate, 130; Na₃GTP, 0.3; HEPES, 10; EGTA, 1; Na-Phosphocreatine, 10 and MgCl₂, 2 (Safari et al., 2017)(Ghaderi et al., 2018). The resistance of these electrodes with the internal solution was 6-8 M Ω . we recorded neuronal membrane potentials in current-clamp mode using the Axopatch 200B amplifier. The sampling rate was 20 kHz, filtered at 2 to 5 kHz and digitized at 10 kHz and transmitted to a computer with NI-DAQ board (PCI-MIO-16E-4, National Instruments). The data were then acquired using custom-made LabVIEW software. The block diagram of our method is mentioned below (Fig. 1).

Even though the intracellular recording technique was blindly done in anesthetized animals and most of the neurons were recorded in superficial layers of the visual cortex, reported in previous studies, in such circumstances, we are much more likely to have recorded regular-spiking putative pyramidal neurons (Liu et al., 2009)(Adesnik, 2017). Although the data reported here is probably a mixture of different types of neurons, they are mainly obtained by pyramidal excitatory neurons, abundant in layers II/III.

2-5- Computational data analysis

We calculated the mean firing rate for every orientation of stimulus (0-360 degrees in steps of 30 degrees) and constructed orientation tuning curve of every neuron before (control condition) and during electrical stimulation of DRN (DBS condition), separately. DBS stands for deep brain stimulation which we use to state the DRN electrical stimulation condition. In the first step, we wanted to know whether there was an effect of DRN stimulation on each recorded neurons' orientation responses. We examined statistically significant changes in the mean firing rate for all trials between control and DBS conditions. According to this, recorded neurons

were classified as unaffected cells, inhibited cells (decreased mean firing rate in DBS condition compared to control condition) and facilitated cells (increased mean firing rate in DBS condition compared to control condition). In the second step, to test how DRN stimulation affected orientation tuning of neurons, we fitted the orientation tuning curve of neurons to the Gaussian function model using the following equation:

$$R(\theta) = Ae^{-\left(\frac{\theta-\theta_0}{\sigma}\right)^2} + R_0$$

Where “ $R(\theta)$ ” is the neuronal response for each oriented stimulus, “ A ” is amplitude or gain; indicates response size at the preferred orientation relative to baseline response, “ θ_0 ” is preferred orientation; the orientation in which the neuron shows the maximum response, “ σ ” is band width reflects the tuning width and also the selectivity of a neuron related to preferred orientation, “ R_0 ” is offset; indicates baseline response of neuron.

Afterwards, the statistical significance of the changes of these four parameters extracted from the model was investigated in the cell groups to determine by what mechanism serotonergic system has affected the changes in orientation tuning curve.

2-6- Statistical analysis

The values were given as Mean $\pm$ SEM; otherwise, they were noted. We used Kolmogorov–Smirnov test for statistical assessment of normality of data distribution. Non-parametric analysis method, Wilcoxon signed-rank test, was used in order to compare the data of two conditions that belong to the same neurons, with or without DRN stimulation.

3- Result

We recorded 63 neurons in layers I and II/III of primary visual cortex using in vivo whole-cell patch-clamp recording before and during electrical stimulation of DRN. Then, each neuron's orientation tuning curve was constructed in control and DBS conditions and their orientation selectivity was examined. 32 of 63 neurons showed orientation selectivity and the others were excluded from more analysis. As mentioned in previous studies, unlike what is seen in monkeys and cats, where most neurons show orientation selectivity, in mice, more than 50% of neurons do not show it (Jeyabalaratnam et al., 2013). Twenty-five neurons showed a significant effect of DRN electrical stimulation on the mean firing rate. 19 of 25 neurons (59.4%) were inhibited and 6 of 25 neurons (18.8%) facilitated by DRN stimulation. There was not any meaningful change in the mean firing rate between control and DBS conditions in 7 of 32 neurons (21.8%).

So, we called them unaffected cells. Therefore, the dominant effect of serotonergic system modulation on recorded neurons' visual response was suppression. The mean firing rate and mean membrane potential (average across the orientation tuning curve) were compared in control and DBS conditions (fig 2). At the population level of inhibited cells, DRN stimulation significantly reduced the mean firing rate in DBS condition ($P = 0.04$, Wilcoxon signed-rank test). Still, there was no meaningful difference in mean membrane potential between two conditions ($P = 0.12$, Wilcoxon signed-rank test).

After fitting the orientation tuning curve of neurons to the Gaussian function model, it is possible to be four different mechanisms according to changes in four parameters of the model (fig. 3a).

1. Changes in selectivity measure can cause the sharpening or broadening of orientation tuning curve; it is correlated to the change in band width (fig. 3b).
2. Changes in preferred orientation (fig. 3c).
3. Changes in baseline response; it is correlated to the change in offset, which can be as an additive (increase in baseline response) or subtractive effect (decrease in baseline response) (fig. 3d).
4. Changes in gain; it is correlated to the change in amplitude, which can be as multiplicative (increase in amplitude) or divisive effect (decrease in amplitude) (Wilson et al., 2012)(Zinke et al., 2006) (fig. 3e).

In our study, the main effect of DRN stimulation on neurons' visual response was suppression, so we focused on inhibited cells. Investigating four parameters showed no remarkable change occurred in preferred orientation during the electrical stimulation of DRN ($P = 0.74$, Wilcoxon signed-rank test) (fig. 4a). We did not find a statistically meaningful difference in band width between control and DBS conditions ($P = 0.87$, Wilcoxon signed-rank test) (fig 4b). Moreover, the offset parameter decreased in all inhibited cells that it was statistically significant. So, the subtractive effect of DRN stimulation was observed on neurons' visual response ($P = 0.0001$, Wilcoxon signed-rank test) (fig. 4c). In Fig. 4d, the scatter plot of amplitude changes has been shown in control and DBS conditions. Amplitude increased in 3 of 19 inhibited cells, and it did not change in 2 of 19 inhibited cells and showed a decrease in the rest of cells. At the population level of inhibited cells, DRN stimulation statistically decreased the gain of visual response of neurons in divisive effect ($P = 0.02$, Wilcoxon signed-rank test).

Examples of different effects of DRN stimulation on orientation tuning curves of inhibited neurons have been shown in Fig. 5. Fig. 5a is a sample for an inhibited cell whose electrical stimulation of DRN caused decreased baseline response in orientation tuning. Therefore, DRN stimulation decreased neurons' visual response in the subtractive effect in DBS condition compared to control. An example of the inhibited cell has been shown in Fig. 5b. The electrical stimulation of DRN led to a decrease in the gain of visual response of cortical neurons in the divisive effect in DBS condition compared to control.

In facilitated neurons by DRN stimulation; statistically, significant changes in model parameters were limited to offset parameter only ($P = 0.03$, Wilcoxon signed-rank test). Because the number of cells in this group was small, more studies are needed.

Finally, we would have like to know whether DRN stimulation affected spontaneous activity or not. For this question, we performed the experiment with a blank stimulus again, and the response of neurons was recorded before and during the electrical stimulation of DRN. We investigated mean membrane potential changes and the mean firing rate of recorded cells ($n=32$). We did not observe any statistically meaningful differences between control and DBS conditions (Fig. 6).

4- Discussion

The principal goal of this research was to investigate the serotonergic modulation of the neural responses to the stimulus orientation in the layers I and II/III of primary visual cortex using the whole-cell patch-clamp recording technique in anesthetized mice. Given that DRN is the main source of serotonergic neuromodulatory neurons in the brain projecting to the visual cortex, we used its electrical stimulation to increase cortical projecting serotonergic neurons' activity. Although the electrical stimulation of DRN is not precisely similar to specific activation of serotonergic neurons, they like other neuromodulator neurons, show highly synchronous locked-in stimulus firing activity (Lottem et al., 2016)(Berkes et al., 2011) and can be used as a model for serotonergic system activation.

Our findings showed that serotonergic system might significantly decrease the mean firing rate by about 60% of the recorded neurons compared to control condition and increased it in a small number of neurons (about 20%). However, the serotonergic system showed a bi-directional effect; its dominant effect was suppression. So, it caused to decrease the visual response of neurons, consistent with others' studies (Waterhouse et al., 1990)(Petzold et al., 2009)(Watakabe et al., 2009)(Seillier et al., 2017)(Azimi et al., 2020)(Michaël et al., 2019).

Investigation of the mechanism by which the serotonergic system decreased visual response revealed that the serotonergic system significantly reduced the baseline response in all inhibited neurons, indicating a subtractive effect. On the other hand, the serotonergic system decreased the gain of visual response through the divisive mechanism in most inhibited neurons. In agreement with our results, Azimi et al. in 2020 using wide-field calcium imaging, multi-unit recording and optogenetic stimulation of DRN reported serotonergic modulation cause two independent inhibitory components including the effect on decreasing gain or amplitude called divisive suppression and the effect on decreasing baseline response called subtractive suppression, each of which is mediated by a specific type of serotonergic receptors.

There was no statistically significant difference in preferred orientation and band width between control and DBS conditions. Therefore, the serotonergic system did not change the preferred orientation and selectivity of the recorded neurons. In 2017, a study by Seillier et al., Which used extracellular recording technique and serotonin iontophoresis in the primary visual cortex of awaking macaque, serotonergic system decreased visual response gain without changing tuning characteristics of neurons including preferred orientation and selectivity.

In our study, there also was not meaningful changes in mean membrane potential and mean firing rate of neurons' spontaneous activity before and during electrical stimulation of DRN, which is in agreement with some previous studies (Michaël et al., 2019). In contrast, some studies reported decreased spontaneous activity in neuronal response by serotonin modulation (Lottem et al., 2016)(Seillier et al., 2017)(Azimi et al., 2020)(Azimi et al., 2018). As Lottem et al. reported in 2016, serotonin reduced neurons' spontaneous activity in the olfactory cortex without altering stimulus-evoked activity. Serotonin cause to change the weight of two aspects of recorded signals in sensory cortex, including spontaneous and stimulus-evoked activities (Azimi et al., 2020)(Azimi et al., 2018). It was suggested that decreasing gain might decrease the priority of a sensory stimulus (Seillier et al., 2017). Otherwise, suppose it only influences the spontaneous response but does not affect the evoked responses (Lottem et al., 2016). In that case, it may enhance the signal-to-noise ratio of a neuron and consequently, increase the stimulus's priority effectively.

Because there are many subtypes of serotonin receptor in the cortex, some of these diverse effects of serotonin probably depend on the postsynaptic compounds of the serotonin receptor subtypes on different neurons in various layers of the cortex (Tseng & Atzori, 2007). Among the subtypes of serotonin receptors, the 5-HT₁ receptors activation leads to an enhance in

potassium ions' conduction throughout the cell membrane, followed by a reduction in excitability of neuron (McCormick et al., 1993). In contrary, 5-HT₂ and 5-HT₄ receptors enhance the spiking activity and excitability of neurons by reducing potassium ions rest conductance and depolarizing the cell membrane (Andrade, 1991)(Panicker et al., 1991)(Bockaert et al., 2010)(Bockaert et al., 2011). Both 5-HT_{1A} and 5-HT_{2A} receptors are major factors in the function of serotonin on the cortical neurons. Studies done by in situ hybridization techniques have shown the simultaneous existence of 5-HT_{2A} and 5-HT_{1A} receptor mRNAs in most pyramidal cells (about 80%) in the cortex. Serotonin can hyperpolarize pyramidal neurons by activating 5-HT_{1A} receptors due to the opening of G protein-coupled inwardly-rectifying potassium channel. This hyperpolarization reduces the spiking activity of excitatory neurons. Serotonin can simultaneously depolarize the same cells via 5-HT_{2A} receptors activation and enhance their excitation. Although there are several hypotheses, it is unclear which elements define the inhibitory or excitatory response to serotonin in a target population of excitatory neurons. At the same way, the effect of these two receptors in modulating role of GABAergic interneuron is unclear. Studies of anatomical bases obviously show a major percentage of interneurons in layers II-VI that have either 5-HT_{2A} receptors or 5-HT_{1A} receptors (Tseng & Atzori, 2007).

Moreover, 5-HT₃ receptors are implicated in controlling the function of GABAergic interneurons. The 5-HT₃ receptors expressing interneurons are located mostly in superficial layers, that play an essential effect in regulating the sensory inputs to the upper parts and tufts of pyramidal neurons's apical dendrites. The function of 5-HT₃ receptors mediates rapid synaptic transmission (Roerig et al., 1997)(Férezou et al., 2002). The 5-HT₃ receptors activation reduces the size of spike and excitation properties in ferret visual cortex, possibly due to increased GABAergic synaptic activity (Roerig & Katz, 1997). Also, the 5-HT_{1B} receptors activation by serotonin have presynaptic modulation of glutamatergic and GABAergic inputs to pyramidal neurons (Tseng & Atzori, 2007)(Xiang & Prince, 2003).

There is a large topographic organization in the dorsal raphe nucleus (Fiser et al., 2010), so other brain's cortical and subcortical areas are also modulated by serotonergic neuromodulatory neurons, affecting activity in the primary visual cortex (Azimi et al., 2018). Moreover, as regards some serotonergic neurons may also use glutamate co-transmission (Ranade & Mainen, 2009)(Muzerelle et al., 2016)(Gilbert & Li, 2013), it can be expected that part of the various effects of serotonin depend on the indirect activity of other circuits (non-

serotonin) and cross talk between the serotonergic system and other neurotransmitters (Tseng & Atzori, 2007).

In overall, the modulatory effect of the serotonergic system on visual sensory processing was, decrease in neural response that can modify the balance between internal and external sensory signals. Perception and sensory cognition is the result of integrating two components (Michaie et al., 2019). As, serotonergic system function may modulate cortical states and influence higher cognition functions such as decision and perception, long-term dysfunction of the serotonergic system may lead to psychiatric disorders including schizophrenia, depression and stress disorder (Azimi et al., 2018). Such studies can be helpful in diagnosis and treatment by providing new methods for rebalancing these components.

Acknowledgments: This article has been extracted from the thesis written by Mrs sareh rostami in School of Medicine Shahid Beheshti University of Medical Sciences (Registration No: 8).

References

- Adesnik, H. (2017). Synaptic Mechanisms of Feature Coding in the Visual Cortex of Awake Mice. *Neuron*, 95(5), 1147-1159.
- Andrade, R. (1991). The electrophysiology of serotonin receptor subtypes. *Serotonin Receptor Subtypes: Basic and Clinical Aspects*, 103–124.
- Aston-Jones, G., & Cohen, J. D. (2005). Adaptive gain and the role of the locus coeruleus–norepinephrine system in optimal performance. *Journal of Comparative Neurology*, 493(1), 99–110.
- Azimi, Z., Barzan, R., Spoida, K., Surdin, T., Wollenweber, P., Mark, M. D., Herlitze, S., & Jancke, D. (2020). Separable gain control of ongoing and evoked activity in the visual cortex by serotonergic input. *ELife*, 9, 1–29.
- Azimi, Z., Spoida, K., Barzan, R., Wollenweber, P., Mark, M., Herlitze, S., & Jancke, D. (2018). Subtraction and division of visual cortical population responses by the serotonergic system. *BioRxiv*, 1–38.
- Berkes, P., Orbán, G., Lengyel, M., & Fiser, J. (2011). Spontaneous cortical activity reveals hallmarks of an optimal internal model of the environment. *Science*, 331(6013), 83–87.
- Bockaert, J., Claeysen, S., Compan, V., & Dumuis, A. (2011). 5-HT₄ receptors, a place in the sun: act two. *Current Opinion in Pharmacology*, 11(1), 87–93.
- Bockaert, J., Perroy, J., Bécamel, C., Marin, P., & Fagni, L. (2010). GPCR interacting proteins (GIPs) in the nervous system: Roles in physiology and pathologies. *Annual Review of Pharmacology and Toxicology*, 50, 89–109.
- Calizo, L. H., Akanwa, A., Ma, X., Pan, Y., Lemos, J. C., Craige, C., Heemstra, L. A., & Beck, S. G. (2011). Raphe serotonin neurons are not homogenous: electrophysiological, morphological and neurochemical evidence. *Neuropharmacology*, 61(3), 524–543.
- Carcea, I., & Froemke, R. C. (2013). Cortical plasticity, excitatory–inhibitory balance, and sensory perception. In *Progress in brain research*, 207, 65–90.
- Cumming, B. G., & Nienborg, H. (2016). Feedforward and feedback sources of choice probability in neural population responses. *Current Opinion in Neurobiology*, 37, 126–132.

- Curzon, G. (1988). Serotonergic mechanisms of depression. *Clinical Neuropharmacology*, 11, S11-20.
- Dahlström, A., & Fuxe, K. (1964). Localization of monoamines in the lower brain stem. *Experientia*, 20(7), 398–399.
- Férezou, I., Cauli, B., Hill, E. L., Rossier, J., Hamel, E., & Lambolez, B. (2002). 5-HT₃ receptors mediate serotonergic fast synaptic excitation of neocortical vasoactive intestinal peptide/cholecystokinin interneurons. *Journal of Neuroscience*, 22(17), 7389–7397.
- Fiser, J., Berkes, P., Orbán, G., & Lengyel, M. (2010). Statistically optimal perception and learning: from behavior to neural representations. *Trends in Cognitive Sciences*, 14(3), 119–130.
- Ghaderi, P., Marateb, H. R., & Safari, M.-S. (2018). Electrophysiological profiling of neocortical neural subtypes: a semi-supervised method applied to in vivo whole-cell patch-clamp data. *Frontiers in Neuroscience*, 12, 823.
- Gilbert, C. D., & Li, W. (2013). Top-down influences on visual processing. *Nature Reviews Neuroscience*, 14(5), 350–363.
- González-Maeso, J., Ang, R. L., Yuen, T., Chan, P., Weisstaub, N. V, López-Giménez, J. F., Zhou, M., Okawa, Y., Callado, L. F., & Milligan, G. (2008). Identification of a serotonin/glutamate receptor complex implicated in psychosis. *Nature*, 452(7183), 93–97.
- Gu, Q. (2002). Neuromodulatory transmitter systems in the cortex and their role in cortical plasticity. *Neuroscience*, 111(4), 815–835.
- Gu, Qiang. (2003). Contribution of acetylcholine to visual cortex plasticity. *Neurobiology of Learning and Memory*, 80(3), 291–301.
- Harris, K. D., & Thiele, A. (2011). Cortical state and attention. *Nature Reviews Neuroscience*, 12(9), 509–523.
- Hornung, J., Fritschy, J., & Törk, I. (1990). Distribution of two morphologically distinct subsets of serotonergic axons in the cerebral cortex of the marmoset. *Journal of Comparative Neurology*, 297(2), 165–181.
- Jacob, S. N., & Nienborg, H. (2018). Monoaminergic neuromodulation of sensory processing. *Frontiers in Neural Circuits*, 12, 51.

- Jeyabalaratnam, J., Bharmauria, V., Bachatene, L., Cattani, S., Angers, A., & Molotchnikoff, S. (2013). Adaptation shifts preferred orientation of tuning curve in the mouse visual cortex. *PloS One*, 8(5), e64294.
- Krnjević, K., & Phillis, J. W. (1963). Iontophoretic studies of neurones in the mammalian cerebral cortex. *The Journal of Physiology*, 165(2), 274.
- Lesch, K.-P., & Waider, J. (2012). Serotonin in the modulation of neural plasticity and networks: implications for neurodevelopmental disorders. *Neuron*, 76(1), 175–191.
- Liu, B. H., Li, P., Li, Y. T., Sun, Y. J., Yanagawa, Y., Obata, K., Zhang, L. I., & Tao, H. W. (2009). Visual receptive field structure of cortical inhibitory neurons revealed by two-photon imaging guided recording. *Journal of Neuroscience*, 29(34), 10520–10532.
- Lottem, E., Lörincz, M. L., & Mainen, Z. F. (2016). Optogenetic activation of dorsal raphe serotonin neurons rapidly inhibits spontaneous but not odor-evoked activity in olfactory cortex. *Journal of Neuroscience*, 36(1), 7–18.
- Lucki, I. (1992). 5-HT₁ receptors and behavior. *Neuroscience & Biobehavioral Reviews*, 16(1), 83–93.
- Lucki, I. (1998). The spectrum of behaviors influenced by serotonin. *Biological Psychiatry*, 44(3), 151–162.
- McCormick, D. A., Wang, Z., & Huguenard, J. (1993). Neurotransmitter control of neocortical neuronal activity and excitability. *Cerebral Cortex*, 3(5), 387–398.
- Michaël, A. M., Parker, P. R. L., & Niell, C. M. (2019). A Hallucinogenic Serotonin-2A Receptor Agonist Reduces Visual Response Gain and Alters Temporal Dynamics in Mouse V1. *Cell Reports*, 26(13), 3475–3483.
- Mokler, D. J., Lariviere, D., Johnson, D. W., Theriault, N. L., Bronzino, J. D., Dixon, M., & Morgane, P. J. (1998). Serotonin neuronal release from dorsal hippocampus following electrical stimulation of the dorsal and median raphe nuclei in conscious rats. *Hippocampus*, 8(3), 262–273.
- Muller, C. P., & Cunningham, K. A. (2020). *Handbook of the behavioral neurobiology of serotonin*. Academic Press.
- Muzerelle, A., Scotto-Lomassese, S., Bernard, J. F., Soiza-Reilly, M., & Gaspar, P. (2016).

- Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5–B9) to the forebrain and brainstem. *Brain Structure and Function*, 221(1), 535–561.
- Ögren, S. O., Eriksson, T. M., Elvander-Tottie, E., D’Addario, C., Ekström, J. C., Svenningsson, P., Meister, B., Kehr, J., & Stiedl, O. (2008). The role of 5-HT_{1A} receptors in learning and memory. *Behavioural Brain Research*, 195(1), 54–77.
- Panicker, M. M., Parker, I., & Miledi, R. (1991). Receptors of the serotonin 1C subtype expressed from cloned DNA mediate the closing of K⁺ membrane channels encoded by brain mRNA. *Proceedings of the National Academy of Sciences*, 88(6), 2560–2562.
- Peroutka, S. J. (1991). *Serotonin receptor subtypes: Basic and clinical aspects*. New York.
- Petzold, G. C., Hagiwara, A., & Murthy, V. N. (2009). Serotonergic modulation of odor input to the mammalian olfactory bulb. *Nature Neuroscience*, 12(6), 784–791.
- Pollak Dorocic, I. (2016). Anatomical and functional characterization of serotonergic neurocircuitry.
- Ranade, S. P., & Mainen, Z. F. (2009). Transient firing of dorsal raphe neurons encodes diverse and specific sensory, motor, and reward events. *Journal of Neurophysiology*, 102(5), 3026–3037.
- Reader, T. A. (1978). The effects of dopamine, noradrenaline and serotonin in the visual cortex of the cat. *Experientia*, 34(12), 1586–1588.
- Roerig, B., & Katz, L. C. (1997). Modulation of Intrinsic Circuits by Serotonin 5-HT₃ Receptors in Developing Ferret Visual Cortex. *Journal of Neuroscience*, 17(21), 8324–8338.
- Roerig, B., Nelson, D. A., & Katz, L. C. (1997). Fast synaptic signaling by nicotinic acetylcholine and serotonin 5-HT₃ receptors in developing visual cortex. *Journal of Neuroscience*, 17(21), 8353–8362.
- Safari, M.-S., Mirnajafi-Zadeh, J., Hioki, H., & Tsumoto, T. (2017). Parvalbumin-expressing interneurons can act solo while somatostatin-expressing interneurons act in chorus in most cases on cortical pyramidal cells. *Scientific Reports*, 7(1), 1–14.
- Seillier, L., Lorenz, C., Kawaguchi, K., Ott, T., Nieder, A., Pourriahi, P., & Nienborg, H.

- (2017). Serotonin decreases the gain of visual responses in awake macaque V1. *Journal of Neuroscience*, 37(47), 11390–11405.
- Tseng, K.-Y., & Atzori, M. (2007). *Monoaminergic modulation of cortical excitability*. Springer.
- Waselus, M., Galvez, J. P., Valentino, R. J., & Van Bockstaele, E. J. (2006). Differential projections of dorsal raphe nucleus neurons to the lateral septum and striatum. *Journal of Chemical Neuroanatomy*, 31(4), 233–242.
- Watakabe, A., Komatsu, Y., Sadakane, O., Shimegi, S., Takahata, T., Higo, N., Tochitani, S., Hashikawa, T., Naito, T., Osaki, H., Sakamoto, H., Okamoto, M., Ishikawa, A., Hara, S. I., Akasaki, T., Sato, H., & Yamamori, T. (2009). Enriched expression of serotonin 1B and 2A receptor genes in macaque visual cortex and their bidirectional modulatory effects on neuronal responses. *Cerebral Cortex*, 19(8), 1915–1928.
- Waterhouse, B. D., Ausim Azizi, S., Burne, R. A., & Woodward, D. J. (1990). Modulation of rat cortical area 17 neuronal responses to moving visual stimuli during norepinephrine and serotonin microiontophoresis. *Brain Research*, 514(2), 276–292.
- William Moreau, A., Amar, M., Le Roux, N., Morel, N., & Fossier, P. (2010). Serotonergic fine-tuning of the excitation–inhibition balance in rat visual cortical networks. *Cerebral Cortex*, 20(2), 456–467.
- Wilson, N. R., Runyan, C. A., Wang, F. L., & Sur, M. (2012). Division and subtraction by distinct cortical inhibitory networks in vivo. *Nature*, 488(7411), 343–348.
- Xiang, Z., & Prince, D. A. (2003). Heterogeneous actions of serotonin on interneurons in rat visual cortex. *Journal of Neurophysiology*, 89(3), 1278–1287.
- Yuen, E. Y., Jiang, Q., Chen, P., Gu, Z., Feng, J., & Yan, Z. (2005). Serotonin 5-HT_{1A} receptors regulate NMDA receptor channels through a microtubule-dependent mechanism. *Journal of Neuroscience*, 25(23), 5488–5501.
- Zinke, W., Roberts, M. J., Guo, K., McDonald, J. S., Robertson, R., & Thiele, A. (2006). Cholinergic modulation of response properties and orientation tuning of neurons in primary visual cortex of anaesthetized Marmoset monkeys. *European Journal of Neuroscience*, 24(1), 314–328.

Legends:

Figure 1. Schematic representation of stereotaxic surgery and whole-cell patch-clamp recording of neurons from the superficial layer of V1 while electrical stimulation of DRN in anesthetized mice presenting drifting grating stimulus.

Figure 2. The comparison of visual evoked activity of recorded neurons before and during electrical stimulation of DRN. a) Mean firing rate and, b) Mean membrane potential of neurons in control and DBS conditions. '+' '*' 'o' represent inhibited, facilitated and unaffected cells, respectively.

Figure 3. Schematic presentation of the Gaussian function model and four predicted curves. a) The orientation tuning curve was presented as a Gaussian function, parameters were explained before. b) mechanism by which selectivity measure changes. c) mechanism by which preferred orientation changes, and there is a shift in tuning curve. d) mechanism by which baseline response changes. Increase and decrease in baseline response lead to additive and subtractive effect, respectively. e) mechanism by which amplitude changes. Increase and decrease in amplitude lead to multiplicative and divisive effect, respectively. The black curve is control, the pink curve is increase, and the blue curve is decrease in response (see arrows).

Figure 4. The scatter plots of four extracted parameters from the model. a) preferred orientation ($P = 0.74$, Wilcoxon signed-rank test), b) band width ($P = 0.87$, Wilcoxon signed-rank test), c) offset; most data points related to inhibited cells drop below the diagonal line, presenting that baseline response was decreased by DRN stimulation ($P = 0.0001$, Wilcoxon signed-rank test). and d) amplitude in control and DBS conditions. Most data points of inhibited cells drop below the diagonal line, presenting which DRN stimulation significantly decreased amplitude in most neurons ($P = 0.02$, Wilcoxon signed-rank test). Diagonal lines present 1 to 1 ratios. '+' '*' 'o' represent inhibited, facilitated and unaffected cells, respectively.

Figure 5. Cell examples of orientation tuning curve before and during electrical stimulation of DRN. a) an inhibited neuron by DRN stimulation that its baseline response was decreased in DBS condition compared to control and it shows subtractive effect. b) an inhibited neuron by DRN stimulation that its amplitude was decreased in DBS condition compared to control and it shows divisive effect. The black curve is control; the pink curve is DBS condition.

Figure 6. The comparison of spontaneous activity in recorded neurons before and during electrical stimulation of DRN a) Mean membrane potential and, b) Mean firing rate of neurons in control and DBS conditions (n=32).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6