Title: Co-Exposure to Toluene and Noise Made Synergistic and Antagonistic Effects on Some Neurotoxic Parameters in New Zealand White Rabbits

Authors: Amirreza Abouee-Mehrizi^{1,2}, Masoud Motalebi Kashani³, Yahya Rasoulzadeh^{4*}, Ahmad Mehdipour⁵, Nafiseh Nasirzadeh⁶, Omid Shatouei-Gharenjeh¹, Abbas Ebrahimi-Kalan⁷

1. *Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran.*
2. *Department of Occupational Health Engineering, Faculty of Health, Tabriz University of Medical Sciences, Tabriz, Iran.*
3. *Occupational Health & Safety Department, Kashan University of Medical Sciences, Kashan, Iran.*
4. *Department of Occupational Health Engineering, Tabriz University of Medical Sciences, Tabriz, Iran.*
5. *Department of Tissue Engineering, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran.*
6. *Department of Occupational Health Engineering, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.*
7. *Neurosciences Research Center (NRC), Department of Neurosciences and Cognitive, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran.*

***Corresponding author:** Yahya Rasoulzadeh, Department of Occupational Health Engineering, Tabriz University of Medical Sciences, Tabriz, Iran. E-mail: rasoulzadehy@tbzmed.ac.ir

To appear in: *Basic and Clinical Neuroscience*

Received date: 2021/03/28

Revised date: 2021/06/02

Accepted date: 2021/10/05

This is a “Just Accepted” manuscript, which has been examined by the peer-review process and has been accepted for publication. A “Just Accepted” manuscript is published online shortly after its acceptance, which is prior to technical editing and formatting and author proofing. *Basic and Clinical Neuroscience* provides “Just Accepted” as an optional and free service which allows authors to make their results available to the research community as soon as possible after acceptance. After a manuscript has been technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as a published article. Please note that technical editing may introduce minor changes to the manuscript text and/or graphics which may affect the content, and all legal disclaimers that apply to the journal pertain.

Please cite this article as:

Abouee-Mehrizi, A., Rasoulzadeh, Y., Mehdipour, A., Nasirzadeh, N., Shatouei-Gharenjeh, O., & Ebrahimi-Kalan, A. (In Press). Co-Exposure to Toluene and Noise Made Synergistic and Antagonistic Effects on Some Neurotoxic Parameters in New Zealand White Rabbits. *Basic and Clinical Neuroscience*. Just Accepted publication Nov. 2, 2022. Doi: <http://dx.doi.org/10.32598/bcn.2022.3301.1>

DOI: <http://dx.doi.org/10.32598/bcn.2022.3301.1>

Abstract

Introduction: Numerous physical and chemical agents can induce destructive effects on the brain tissue. Noise and toluene, which are some of these harmful agents, have significant adverse effects on the brain tissue. This work aims to investigate the neurotoxic changes induced by co-exposure to toluene and noise.

Methods: Totally, 24 male white New Zealand rabbits were randomly segregated into 4 groups including toluene exposure, noise exposure, co-exposure to noise and toluene, and control. This in-vivo study tested the neurotoxic effects of exposure to 1000 ppm toluene and 100 dB noise during two weeks (8 h/day). The serum levels of brain-derived neurotrophic factor- α (BDNF- α) and malondialdehyde (MDA), total antioxidant capacity (TAC), glutathione peroxidase (GPx), superoxide dismutase (SOD), and catalase levels in the brain tissue were measured. Moreover, hematoxylin and eosin staining was utilized for brain pathological analysis.

Results: Exposure to noise increased TAC level in the cerebral cortex. Co- exposure to toluene and noise increased the serum level of BDNF- α . Nevertheless, exposure to noise decreased the level of BDNF- α in serum. On the other hand, hispathological examinations using hematoxylin-eosin (H&E) exhibited that different signs of inflammation such as lymphocyte infiltration, pyknosis, vacuolization, and chromatolysis were induced by exposure to noise and toluene in the cerebellum, hippocampus, and frontal section in the brain tissue. In addition, simultaneous exposure to toluene and noise induced antagonistic and synergistic changes on some neurotoxic parameters.

Conclusion: Exposure to noise and toluene, which caused inflammation in the brain tissue cells, could be a noticeable risk factor for neurological system.

Keywords: Noise, Toluene, Brain, Neurotoxicity, Oxidative stress

1. Introduction

Noise is a fundamental property of different activities which is being produced by changes in air pressure (Snyder, 2000). Excessive noise hurts various systems in the body including the cardiovascular system, auditory system, psychological system, and neural system (Babisch, 2003; Brunoni et al., 2008). It has been proven that exposure to noise causes some disorders in the brain tissue such as congestion, edema, blood vessel expansion, neuroglial hyperplasia, and mild atypia (Xue et al., 2014). Regrettably, the main cause or intensifying factors of many disasters and accidents which could be related to the neurological and psychological disorders provoked by exposure to noise such as enhancing the occurrence of violence and increasing workplace accidents (Toppila et al., 2009) may not be well detected or resolved.

Toluene is one of the most suitable solvents that has various usages in numerous industries such as plastics, rubber, leather, adhesives, and pharmaceuticals (Acton, 2013). Previous studies have demonstrated that exposure to toluene induces different diseases in various organs of the body, including the liver and kidneys (Schardein and Macina, 2006). The psychological and neurotoxic effects of toluene are complicated (Ruiz et al., 2011) and include loss of balance, headache, weakness, fatigue, and joy feelings. In addition, toluene has a devastating effect on the nervous tissue of the brain (Dobbs, 2009).

There are many different physical and chemical stressors in the industry. Noise and toluene are some of the prevalent physical and chemical stressors in different workplaces. Previous studies have indicated that different stressors could deteriorate some brain disorders (Demir et al., 2017; Xiu et al., 2009).

The debate about health effects induced by Co-exposure to chemical and physical pollutants has gained fresh prominence with many arguing on the manner of the interactions between different chemical and physical harmful factors (Sexton and Linder, 2011; Streffer et al., 2013). As stated by numerous reports and studies, many people in various occupations are not only exposed to noise (such as constructions, factories, nightclubs, and airports) (Tripathy, 2008) or only toluene (such as petrochemical, plastics factories, adhesive factories, and pharmaceutical factories) (Acton, 2013; Miller, 2002), but also simultaneously exposed to noise and toluene (Dobie, 2015; Rea and Patel, 2014). This indicates a need to understand various perceptions of neurotoxicity induced by simultaneous exposure to noise and toluene. Therefore, this experimental study

provides one of the first investigations on how co-exposure to toluene and noise changes biochemical factors and brain tissue in rabbits.

2. Materials and Methods

2.1. Animals and divided groups

Twenty four healthy adult male white New Zealand rabbits (Pasteur Institute of Tehran, Iran) weighing 2.83 ± 0.41 kg were used in this experiment. Animals were kept on a 12:12 hour light/dark circulation period with controlled ambient temperature at $21 \pm 2^\circ\text{C}$ and humidity between 50 to 70% while having access to food (standard rabbit pellet, Behparvar Feed Manufacturing Company, Tehran) and treated water. Animals were housed in the laboratory for 14 days before the experiments to adapt to the environmental conditions of the laboratory. They were randomly segregated into 4 groups ($n=6$ per group) including control group (group 4), noise exposure group (group 1), toluene exposure group (group 2), and Co-exposure to toluene and noise group (group 3). Exposure duration was 14 consecutive days (8 h/day from 9:00 a.m. to 5:00 p.m.) for each group (as well as control group). First of all, control group was placed into the exposure chamber with clean air flowing (toluene = 0 ppm) and < 50 dBA noise (background noise) the exposure duration (14 consecutive days). Then, groups 1 placed in the exposure chamber to exposure to 100 ± 5 decibel (dBA) noise for the exposure duration. After that, group 2 was placed in the same exposure to exposure to 1000 ± 50 parts per million (ppm) for the exposure duration. After all, group 3 was placed in the exposure chamber and exposed to noise and toluene with the same specifications as groups 1 and 2 for the exposure duration. The temperature was $23 \pm 3^\circ\text{C}$ and the relative humidity was 65-80% inside the exposure chamber for all the experimental groups.

2.2. Exposure chamber

Animals in all the groups ($n=6$) were exposed in the same exposure chamber with the dimensions of $50 \times 60 \times 90$ cm³ and made of clear and transparent polycarbonate sheets, which was coated with transparent PET (polyethylene terephthalate) adhesive tape. Specifications of the exposure chamber were chosen according to the animals' welfare requirements (Gad, 2006) and based on the reverberant chamber features (Cobo et al., 2009; Moreno et al., 2000) to generate the same noise specifications inside the exposure chamber and to make noise intensity independent from distance in the chamber.

2.3. Noise exposure set-up

White noise at 100 ± 5 dBA (70-20000 Hz) was produced by Windows noise generator software (Audacity® Software 1.3.12 Beta) and monitored by another Windows software (Cool Edit Pro 2.1 ©1992-2003 Syntrillium Software Corporation). The selected noise was amplified by a power amplifier (3030 W, MULTI TONE) and transported to a loudspeaker. The loudspeaker was placed directly above the chamber on the chamber roof. Noise level was continuously measured using a sound level analyzer (TES 1358 Sound Analyzer Real-Time 1/1 & 1/3 octave band Analyzer) in 1/3 octave band during the exposure (groups 1 and 3). The sound level meter microphone was placed at the same level as the ears of the animals by checking the designed outlets (made of rubber tape and coated with PET adhesive tape) every 30 min in order to monitor noise specifications.

2.4. Toluene exposure set-up

Exposure to 1000 ± 50 ppm toluene happened through its surface evaporation. In summary, constant generation of toluene vapors was done using an impinger (capacity of 250 ml) with content of 100 ml toluene liquid (Merck KGaA, extra pure, Darmstadt, Germany) and via injecting 20 ml toluene into the impinger every 90 min on the periodical basis. The toluene concentration was analyzed during pilot studies using a calibrated humidity-resistant real-time instrument (Phocheck, Ion Science Ltd, Cambs UK 07-01782) and also charcoal absorbents with gas chromatography (Model Agilent 7890A GC System) established on the 1501 NIOSH method. Accordingly, changing the toluene concentration was negligible (5%) during the exposure time in the pilot and real experiments. According to the pilot experiments, the pure air flow rate was 30 L/min and the toluene vapor-contained air flow rate was 3 L/min. Moreover, before starting real exposures, the purity of the toluene vapors was confirmed by air sampling from the exposure chamber (active charcoal samples) and using a gas chromatography-mass spectrometry instrument (Agilent 6890 / 5973 GCMS). To obtain the target concentration of toluene constantly and continuously, a mixer chamber which had the same materials and specifications as the exposure chamber with the dimensions of $50 \times 50 \times 20$ cm³, was used before entering the air into the exposure chamber. To monitor and check toluene concentration, the same real-time instrument (Phocheck) which had been already calibrated and utilized in pilot experiments was used. Air samples were prepared from inside the exposure chamber through

checking outlets designed on four sides in the surrounded walls of the exposure chamber (checking outlets).

2.5. Blood sampling

Blood samples were collected from the marginal vein of the ears of each animal in 5 stages, as mentioned in Fig. 1.

2.6. Tissue sampling and processing

Rabbits were sacrificed after anesthesia by 35 mg/kg ketamine and 5 mg/kg xylazine (IM injection). The animals were sacrificed 14 days after the end of the exposure in each group. The whole brain consisting of cerebral cortex, hippocampus, and cerebellum was dissected out immediately after the animals were scarified and fixed at 10% formalin solution (formaldehyde solution, Merck KGaA, Darmstadt, Germany) at pH 7.2. After the fixation, paraffin blocks were prepared. Moreover, for biochemical analyses, the cerebral cortex of the brain tissues was immediately collected and kept at -80°C until the time of biochemical experiments. Dehydration and paraffin impregnation phases of the tissue samples were made by an autotechnicon device. Then, paraffin molding was carried out and the slides were prepared using microtome with the typical diameter of 5 μ . Eventually, the slides were stained with hematoxylin-eosin (H&E) and histopathologic changes were assessed by light microscopy (Nikon Eclipse E100, Japan).

2.7. Homogeneity of brain tissue

First of all, 100 gr of the cerebral cortex of the brain samples was weighed out of all the groups to measure the oxidant/antioxidant parameters of the brain tissue. Afterwards, they were dissolved in 1 cc PBS buffer and homogenized by a homogenizer device (ultrasonic homogenizer Hielscher UIP-1500HD). Then, the samples were centrifuged by a centrifuge device at 1000 g at 4°C for 20 min. Required samples were taken from the clear supernatant liquid of the centrifuged samples using a sampler and placed in microtubes. At that time, the supernatant protein liquid of the cerebral cortex was measured using a commercial kit (Pars Azmoon Company, Karaj, Iran). Finally, the parameters in the supernatant liquid were normalized by the measured amount of supernatant protein liquid of the brain tissue and valued per mg of tissue protein.

2.8. Determining oxidant and anti-oxidant factors in the cerebral cortex using spectrophotometric method

Total antioxidant capacity (TAC) was measured by a Naxifer™ diagnosis kit (total antioxidant capacity assay kit, Navand Lab Kit, Urmia, Iran). Level of glutathione peroxidase (GPx) was measured by a Biorexfars diagnosis kit (glutathione peroxidase kit, production code BXC0551, Shiraz, Iran). Catalase level was measured by a Cayman chemical company kit (catalase assay kit, Item no. 707002) and superoxide dismutase (SOD) level was measured by a Nasdox™ diagnosis kit (non-enzymatic method kit, Navand Lab Kit, Urmia, Iran). All the measurements were done according to the instructions provided by the kit manufacturers; in addition, malondialdehyde (MDA) level was measured using thiobarbituric acid (TBA) method (Janero, 1990; Stewart and Bewley, 1980).

2.9. Determining the serum level of brain-derived neurotrophic factor- α (BDNF- α) using ELISA method

BDNF- α was measured by Rabbit ELISA kits (HANGZHOU EASTBIOPHARM CO., LTD, Hangzhou, China) and based on the manufacturer's instructions (Cat. No: CK-E91855). In this experiment, the State Fax 2100 ELISA plate reader (Awareness Technology, Inc., USA) and the State Fax 2600 ELISA plate washer (Awareness Technology, Inc., USA) were employed.

2.10. Statistical analysis

The GEE (generalized estimating equations) statistical method was used for BDNF- α , so that differences between different repeats in different groups were compared in SPSS software (version 25.0.0.0, Armonk, NY: IBM Corp). Moreover, ANOVA (analysis of variance) statistical method followed by Tukey's pairwise comparisons was used to compare the oxidant and anti-oxidant parameters and the mean of the brain tissue weight/body weight in different groups in Minitab software (*Version 18*). The statistical significance level was considered at 0.05. The logarithm of abnormal parameters was used for doing GEE statistical analysis.

3. Results

3.1. Biochemical changes

According to Fig. 2-E, TAC level in group 1 was significantly higher than that in groups 2, 3, and 4. Figures 2-A, 2-B, 2-C, and 2-D show that catalase, GPx, SOD, and MDA levels in group 1.

According to the findings obtained from this study, BDNF- α level in group 1 was significantly lower than that in group 2 at stages 4 and 5. BDNF- α level in group 1 was significantly lower than that in control group at stage 5. Moreover, BDNF- α level in group 2 significantly decreased at stages 2 and 3 compared to stages 1, 4, and 5. Furthermore, BDNF- α in group 3 was significantly higher than that in groups 1 and 2 at stage 3 and significantly lower than group 2 at stages 4 and 5. BDNF- α in group 3 significantly increased at stage 3 compared to stages 1, 2, and 4. BDNF- α in group 3 significantly increased at stage 5 compared to stage 1 (Fig. 3).

3.2. Brain tissue weight changes

According to Fig. 4, the average brain weight/body weight in noise exposure group was significantly lower than that in the toluene exposure group.

3.3. Histopathological changes

The cerebellum was in normal conditions in control group. However, the population of Purkinje cells decreased in the cerebellum in group 1; in addition, pyknosis appeared in Purkinje cells in this group. The relative reduction in the number of cells was visible in the granular layer in groups 1 and 2. Granular cells not only reduced relatively, but also appeared massively in group 2. Pathological changes were more specific in the cerebellum area in group 3. Reduction in the population of Purkinje cells, decrease in the layer thickness of the granular layer, and reduced granular cells were some of the noticeable changes observed in the cerebellum in group 3 (Fig. 5).

The cortex area in the frontal part of the brain was normal in control group. Nevertheless, gliosis was detected in the prefrontal cortex area in groups 1, 2, and 3 compared to control group. The number of pyknotic cells in groups 1 and 3 appeared higher than group 2 in the prefrontal cortex area. Minor lymphocyte infiltration, which was possibly induced by retrograde effect of toluene and transmitted to the frontal part of the brain cortex through olfactory nerves, was visible in the prefrontal cortex area in group 2. Some pathological changes were noticed in the prefrontal cortex area in group 3 such as disorganization in the cortex layer, relative reduction of pyramidal cells, vacuolization, and chromatolysis (Fig. 6).

The light micrographs of the hippocampus area were normal in control group. Dark cells were reduced in population and also were pyknotic in the hippocampus area in group 1, even more

severely than group 2. However, it was not remarkable compared to group 3. Moreover, the pyramidal cells in group 1 noticeably decreased in population compared to group 2 in the hippocampus area. But it was not remarkable compared to group 3. All the changes reported for group 1 also appeared in group 2 in the hippocampus area, but were milder than in group 3, so that the pathological changes appearing in group 3 were more severe than those in groups 1 and 2 (Fig. 7).

4. Discussion

The results of this study showed that noise exhibited an increase in TAC level in the cerebral cortex. In one study on the effects of industrial noise stress in workers, the serum levels of catalase and TAC were significantly higher in workers exposed to noise stress (Hosseinabadi et al., 2019). Increased SOD level was reported by exposure to noise at 100 dB in the cerebral cortex of the rats' brain (Samson et al., 2007); however, it is possible that increased anti-oxidant factors after exposure to high level of noise is due to induced oxidative stress during exposure to noise. Moreover, activating protective mechanisms in the body such as producing TAC, GPx, and SOD catalase as anti-oxidant factors and decreasing MDA as an oxidative factor are predictable during oxidative stress to prevent the excessive generation of free radicals (Rosenbaum et al., 1994). Furthermore, some of the previous studies have shown that toluene induces oxidative stress in several brain regions by producing reactive oxygen species (ROS) (Kodavanti et al., 2011). As a result, ROS can induce apoptosis in the brain (Redza-Dutordoir and Averill-Bates, 2016). Therefore, co-exposure to toluene and noise changed oxidant and anti-oxidant parameters in the brain tissue.

Brain-derived neurotrophic factor is a protein encoded by a gene called BDNF. This protein promotes the growth and development of the central and peripheral nervous system. It also triggers nerve synapses and establishes neural connections. It is most active in the hippocampus and the upper part of the brain. The highest production of BDNF is in the brain, the amount of which varies in brain damage, especially with environmental stress.

The current study revealed that co-exposure to toluene and noise showed an increase in the serum level of BDNF; however, exposure to noise decreased BDNF serum level compared to the control group. In the study on the effects of inhalation exposure to 5 ppm toluene for 5 days, a decrease in the relative mRNA expression of BDNF, and an increase in the mRNA expression

levels of NF- κ B were reported in the hippocampus of the infant mice for the period of brain development (Win-Shwe et al., 2012). In the study concerning the effects of chronic stress on BDNF level, inducing 80 dB noise stress for 5 weeks made a significant reduction in hippocampal BDNF level in rats (Taliaz et al., 2011). Moreover, another study demonstrated that acute stress caused oxidative stress and, consequently, produced an increase in BDNF level (Hacioglu et al., 2016). It means there is a direct relationship between BDNF level and oxidative stress indicators (Hacioglu et al., 2016). On the other hand, previous studies have shown that some stressors such as noise initiate an increase in corticosterone level (Laugero and Moberg, 2000; Lehmann et al., 2002; Şahin and Gümüşlü, 2007). Consequently, corticosterone can induce a decrease in BDNF level (Jacobsen and Mork, 2006; Schaaf et al., 2000).

This study provides considerable insight into the combined effects of noise and toluene on neurological disorders. Based on the results of the present study related to oxidative stress induced after co-exposure to toluene and noise and according to the multiple combined stressors (Piggott et al., 2015), different synergistic and antagonistic effects induced by exposure to noise and toluene are provided in Appendix 1 and Appendix 2.

This study demonstrated that co-exposure to noise and toluene can induce different histopathological changes in the brain tissue such as lymphocyte infiltration, relative reduction of pyramidal cells, chromatolysis, and vacuolization. There were remarkable signs of inflammation in co-exposure group, even more severely than noise and toluene exposure groups. Based on the results obtained, it seems that even though there were no significant changes induced by noise exposure, it was found some changes in the brain tissue, which if persisted it could be an important alarm. Induced mild atypia, edema, hyperemia, expansion of blood vessels, and glial cell hyperplasia were reported in several areas in the brain tissue in female rats by exposure to noise at 95 dB (4000 Hz) for 2 weeks (Xue et al., 2014). Makhlof et al. (2014) reported a decrease in the thickness of pyramidal cell layer of CA1 (cornu ammonis area 1), CA3 (cornu ammonis area 3), and also the granular cell which exhibited degeneration and loss of many cells after exposure to 100 dB noise during 4 weeks (4h/day) in male albino rats (Makhlof et al., 2014). Abousetta et al. (2014) indicated that exposure to 100 dB white noise for 6 h/day during 4 consecutive weeks induced a decrease in thickness of dentate gyrus (DG) granular cell layer and

pyramidal cell layer with the loss and degeneration of numerous cells in male albino rats (Abousetta et al., 2014).

Enhanced glial fibrillary acidic protein (GFAP)-immunoreactive cytoplasm was demonstrated in the white matter of the cerebellum and in the dentate gyrus of the hippocampus by inhaling 1500 ppm toluene for 4 and 7 days in the brain tissue of the rats (Gotohda et al., 2000). In addition, Gotohda et al. (2000) showed that chronic exposure to 3000 ppm toluene for 12 weeks induced severe degenerative changes, widely dark picnotic nuclei, and shrunken cytoplasm in neurons of different regions of the hippocampus in rats (Gotohda et al., 2000). Kanter (2007) demonstrated that exposure to 3000 ppm toluene for 12 weeks (8 h/day) induced dilated cisternae of endoplasmic reticulum, shrunken cytoplasm, evidently inflamed mitochondria with deteriorated cristae in the hippocampus in male Wistar albino rats (Kanter, 2008). Tas et al. (2012) reported that exposure to the high dose of toluene (5200 mg/kg/gavage) for 2 weeks caused a significant increase in the immune reactivity of Bax in the cerebellum and the brain cortex (Tas et al., 2012). Demir et al. (2017) demonstrated that acute exposure to toluene by a single dose (876 mg/kg) intraperitoneal (IP) injection in the New Zealand rabbits brain caused significantly extreme expansion of the blood vessels, dispersed cell borders, vacuolar degeneration, severe degeneration of the compensation, perivascular demyelination, gliosis, and several necrosis and pyknotic cells (Demir et al., 2017). Moreover, a noticeable enhance in the count of apoptotic cells in the cerebral granule layer was indicated by inhaling 1200-1800 ppm toluene for 6 h per day in rats (Dalgaard et al., 2001). Nevertheless, Seo et al. (2009) indicated that exposure to toluene at the dose of 500 mg/kg did not cause apoptosis in the hippocampus in mice (Seo et al., 2010). Considering the findings proven by previous studies and the results obtained from this study, noise and toluene are some of the noticeable neurotoxic pollutants.

5. Limitations

Although this study improved non-auditory changes induced by exposure to toluene and noise, neurotoxic effects induced by chronic exposure to noise and toluene were not identified. Furthermore, this study did not evaluate other biological and pathological factors during or after exposure. Therefore, further studies are necessary in order to enhance the results obtained from this work.

6. Conclusion

This study indicated that co-exposure to toluene and noise might induce neurotoxicity in white New Zealand rabbits. Histopathological experiments indicated that co-exposure to toluene and noise could aggravate neurotoxic effects. Co-exposure to noise and toluene induced some pathological changes in cerebellum, hippocampus, and frontal section. Nonetheless, further studies are needed to determine the definite neurotoxic effects induced by co-exposure to toluene and noise.

Declaration of interest

There is no conflict of interest.

Accepted Manuscript (Uncorrected Proof)

References

- Abousetta, A., Makhoulf, N.A., El-Beshbishy, R.A. (2014). The effects of concomitant Ginkgo intake on noise induced Hippocampus injury. Possible auditory clinical correlate. Egyptian Journal of Ear, Nose, Throat and Allied Sciences 15(3):231-239.
- Acton, Q.A. (2013). Advances in Toluene Research and Application: 2013 Edition: ScholarlyBrief. ScholarlyEditions, Atlanta, Georgia, USA.
- Babisch, W. (2003). Stress hormones in the research on cardiovascular effects of noise. Noise and health 5(18):1.
- Brunoni, A.R., Lopes, M., Fregni, F. (2008). A systematic review and meta-analysis of clinical studies on major depression and BDNF levels: implications for the role of neuroplasticity in depression. International Journal of Neuropsychopharmacology 11(8):1169-1180.
- Cobo, P., Murillo-Cuesta, S., Cediell, R., Moreno, A., Lorenzo-García, P., Varela-Nieto, I. (2009). Design of a reverberant chamber for noise exposure experiments with small animals. Applied Acoustics 70(8):1034-1040 doi:<https://doi.org/10.1016/j.apacoust.2009.03.005>.
- Dalgaard, M., Hossaini, A., Hougaard, K., Hass, U., Ladefoged, O. (2001). Developmental toxicity of toluene in male rats: effects on semen quality, testis morphology, and apoptotic neurodegeneration. Archives of toxicology 75(2):103-109.
- Demir, M., Mustafa, C., Nadire, E., Atila, Y., Turgay, S. (2017). Effects of acute toluene toxicity on different regions of rabbit brain. Analytical Cellular Pathology. 2017:2805370. doi: 10.1155/2017/2805370
- Dobbs, M.R. (2009). Clinical Neurotoxicology E-Book: Syndromes, Substances, Environments. Elsevier Health Sciences, Kentucky, USA.
- Dobie, R.A. (2015). Medical-Legal Evaluation of Hearing Loss, Third Edition. Plural Publishing, Incorporated, California.
- Gad, S.C. (2006). Animal models in toxicology, Second Edition. Taylor & Francis, Oxfordshire United Kingdom.

- Gotohda, T., Tokunaga, I., Kubo, S.i., Morita, K., Kitamura, O., Eguchi, A. (2000). Effect of toluene inhalation on astrocytes and neurotrophic factor in rat brain. *Forensic science international* 113(1-3):233-238.
- Hacioglu, G., Senturk, A., Ince, I., Alver, A. (2016). Assessment of oxidative stress parameters of brain-derived neurotrophic factor heterozygous mice in acute stress model. *Iran J Basic Med Sci* 19(4):388-393.
- Hosseiniabadi, M.B., Khanjani, N., Ebrahimi, M.H., Mirbadie, S.R., Biganeh, J. (2019). The effects of industrial noise exposure on lipid peroxidation and antioxidant enzymes among workers. *International Archives of Occupational and Environmental Health*:1-6.
- IARC (2020). IARC Monographs on the Identification of Carcinogenic Hazards to Humans. World Health Organization, International Agency for Research on Cancer, Volume 1-125. <https://monographs.iarc.fr/list-of-classifications>.
- Jacobsen. J.P., Mork, A. (2006). Chronic corticosterone decreases brain-derived neurotrophic factor (BDNF) mRNA and protein in the hippocampus, but not in the frontal cortex, of the rat. *Brain Res* 1110(1):221-5 doi:10.1016/j.brainres.2006.06.077.
- Janero, D.R. (1990). Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. *Free Radical Biology and Medicine* 9(6):515-540. doi:[https://doi.org/10.1016/0891-5849\(90\)90131-2](https://doi.org/10.1016/0891-5849(90)90131-2).
- Kanter, M. (2008). *Nigella sativa* and derived thymoquinone prevents hippocampal neurodegeneration after chronic toluene exposure in rats. *Neurochemical research* 33(3):579.
- Kodavanti, P.R.S., Royland, J.E., Richards, J.E., Besas, J., MacPhail, R.C. (2011). Toluene effects on oxidative stress in brain regions of young-adult, middle-age, and senescent Brown Norway rats. *Toxicology and applied pharmacology* 256(3):386-398.
- Laugero, K.D., Moberg, G.P. (2000). Energetic response to repeated restraint stress in rapidly growing mice. *American Journal of Physiology-Endocrinology and Metabolism* 279(1):E33-E43.
- Lehmann, J., Russig, H., Feldon, J., Pryce, C.R. (2002). Effect of a single maternal separation at different pup ages on the corticosterone stress response in adult and aged rats. *Pharmacology Biochemistry and Behavior* 73(1):141-145.

- Makhlouf, N.A., El-Beshbishy, R.A., Abousetta, A. (2014). Ginkgo modulates noise-induced hippocampal damage in male albino rats: a light and electron microscopic study. *Egyptian Journal of Histology* 37(1):159-174.
- Miller, R.L. (2002). *The Encyclopedia of Addictive Drugs*. Greenwood Press, Westport, USA.
- Moreno, A., Ruiz, J., Colina, C.d.l. (2000). Re-visiting Bolt's criterion for homogeneous distribution of normal frequencies in rectangular enclosures. *ACUSTICA*, Madrid, Spain.
- Piggott, J.J., Townsend, C.R., Matthaei, C.D. (2015). Reconceptualizing synergism and antagonism among multiple stressors. *Ecology and evolution* 5(7):1538-1547.
- Rea, W.J., Patel, K.D. (2014). *Reversibility of Chronic Disease and Hypersensitivity, Volume 2: The Effects of Environmental Pollutants on the Organ System*. CRC Press, New York, USA.
- Redza-Dutordoir, M., Averill-Bates, D.A. (2016). Activation of apoptosis signalling pathways by reactive oxygen species. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1863(12):2977-2992 doi:<https://doi.org/10.1016/j.bbamcr.2016.09.012>.
- Rosenbaum, D.M., Kalberg, J., Kessler, J.A. (1994). Superoxide dismutase ameliorates neuronal death from hypoxia in culture. *Stroke* 25(4):857-862.
- Ruiz, P., Strain, E.C., Lowinson, J.H. (2011). *Lowinson and Ruiz's Substance Abuse: A Comprehensive Textbook*. Wolters Kluwer Health/Lippincott Williams & Wilkins.
- Şahin, E., Gümtüçlü, S. (2007). Immobilization stress in rat tissues: alterations in protein oxidation, lipid peroxidation and antioxidant defense system. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 144(4):342-347.
- Samson, J., Sheeladevi, R., Ravindran, R. (2007). Oxidative stress in brain and antioxidant activity of *Ocimum sanctum* in noise exposure. *Neurotoxicology* 28(3):679-685.
- Schaaf, M.J.M., De Kloet, E.R., Vreugdenhil, E. (2000). Corticosterone Effects on BDNF Expression in the Hippocampus Implications for Memory Formation. *Stress* 3(3):201-208. doi:10.3109/10253890009001124.
- Schardein, J.L., Macina, O.T. (2006). *Human Developmental Toxicants: Aspects of Toxicology and Chemistry*. CRC Press.

- Seo, H.S., Yang, M., Song, M.S., Kim, J.S., Kim, S.H., Kim, J.C., Kim, H., Shin, T., Wang, H. and Moon, C. (2010). Toluene inhibits hippocampal neurogenesis in adult mice. *Pharmacology Biochemistry and Behavior* 94(4):588-594.
- Sexton, K., Linder, S.H. (2011). Cumulative risk assessment for combined health effects from chemical and nonchemical stressors. *American Journal of Public Health* 101(S1):S81-S88.
- Snyder, S. D. (2000). *Active Noise Control Primer*. Springer, New York.
- Stewart, R.R., Bewley, J.D. (1980). Lipid peroxidation associated with accelerated aging of soybean axes. *Plant physiology* 65(2):245-248.
- Streffer, C., Bücker, J., Cansier, A., Cansier, D., Gethmann, C.F., Guderian, R., Hanekamp, G., Henschler, D., Pösch, G., Rehbinder, E. and Renn, O. (2013). *Environmental Standards: Combined Exposures and Their Effects on Human Beings and Their Environment*. Springer Berlin Heidelberg.
- Taliaz, D., Loya, A., Gersner, R., Haramati, S., Chen, A., Zangen, A. (2011). Resilience to Chronic Stress Is Mediated by Hippocampal Brain-Derived Neurotrophic Factor. *The Journal of Neuroscience* 31(12):4475 doi:10.1523/JNEUROSCI.5725-10.2011.
- Tas, U., Ayan, M., Kuloglu, T., Suren, M., Cakil, D., Ozyurt, B., Bekar, D.A. and Sarsilmaz, M. (2012). Examination of Apoptotic Effects of High-Dose Toluene on the Brain Cortex and Cerebellum Tissue During the Acute Phase: an experimental study. *European Journal of General Medicine* 9(4).
- Toppila, E., Pyykkö, I., Pääkkönen, R. (2009). Evaluation of the increased accident risk from workplace noise. *International journal of occupational safety and ergonomics* 15(2):155-162.
- Tripathy, D.P. (2008). *Noise Pollution*. A.P.H. Publishing Corporation, New Delhi, India.
- Win-Shwe, T.T., Kunugita, N., Yoshida, Y., Nakajima, D., Tsukahara, S., Fujimaki, H. (2012). Differential mRNA expression of neuroimmune markers in the hippocampus of infant mice following toluene exposure during brain developmental period. *Journal of Applied Toxicology* 32(2):126-134.

Xiu, M.H., Hui, L., Dang, Y.F., De Hou, T., Zhang, C.X., Zheng, Y.L., Kosten, T.R. and Zhang, X.Y. (2009). Decreased serum BDNF levels in chronic institutionalized schizophrenia on long-term treatment with typical and atypical antipsychotics. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 33(8):1508-1512.

Xue, L., Zhang, D., Wang, T., Shou, X. (2014). Effects of high frequency noise on female rat's multi-organ histology. *Noise and Health* 16(71):213.

Accepted Manuscript (Uncorrected Proof)

Fig. 1. Stages of blood sampling. The blood samples collected immediately before exposure and at different moments after the end of exposure for each group.

Fig. 2. Different oxidant and anti-oxidant parameters in the brain tissue; Factors were measured fourteen days subsequently the end of exposure to 100 ± 5 decibel (dB) noise (Noise, $n=6$), 1000 ± 50 parts per million (ppm) toluene (Toluene, $n=6$), co-exposure to 100 ± 5 dB noise and 1000 ± 50 ppm toluene (Noise+Toluene, $n=6$), and control group (Control, $n=6$). Bars represent the arithmetic mean and standard error (mean $\pm$ SE). The asterisk (*) exhibits significant differences between noise and other groups (P -value < 0.001). GPx is glutathione peroxidase. MDA is malondialdehyde. SOD is superoxide dismutase. TAC is total antioxidant capacity. Exposure routes: Inhalation exposure to toluene and hearing exposure to noise

Fig. 3. Serum level of brain-derived neurotrophic factor- α (BDNF- α) in exposure groups in different stages; Points represent the arithmetic mean and standard error (mean $\pm$ SE). Groups include control group (group 4, n=6), exposure to noise at 100 ± 5 decibel (dB) (group 1, n=6), exposure to toluene at 1000 ± 50 parts per million (ppm) (group 2, n=6), and co-exposure to 100 ± 5 noise and 1000 ± 50 ppm toluene (group 3, n=6). Stages include pre-exposure (stage 1), immediately after the end of exposure (stage 2), 3 days after the end of exposure (stage 3), 7 days after the end of exposure (stage 4), and 14 days after the end of exposure (stage 5). Exposure to noise significantly decreased BDNF- α compared to control and toluene exposure groups on the 14th day after exposure. Moreover, co-exposure to noise and toluene significantly increased BDNF- α compared to noise and toluene exposure groups on the 3rd day after exposure. Furthermore, co-exposure to noise and toluene significantly decreased BDNF- α compared to toluene exposure group on the 7th and 14th days after exposure. Exposure routes: Inhalation exposure to toluene and hearing exposure to noise

Fig. 4. Geometric mean of brain weight/body weight in different groups; The parameter was measured fourteen days after the end of exposure to 100 ± 5 decibel (dB) noise (Noise, $n=6$), 1000 ± 50 parts per million (ppm) toluene (Toluene, $n=6$), co-exposure to 100 ± 5 dB noise and 1000 ± 50 ppm toluene (Noise+Toluene, $n=6$), and control group ($n=6$). Bars represent the geometric mean and standard error (mean $\pm$ SE). The asterisks (*) exhibit significant differences between noise and toluene exposure groups (P-value <0.05). Exposure routes: Inhalation exposure to toluene and hearing exposure to noise

Fig. 5. Light micrograph of cerebellum lesions stained with hematoxylin and eosin (H&E) at 10x (A, B, C, and D) and (40x a, b, c, and d); Lesions were removed fourteen days later the end of exposure to 100 ± 5 decibel (dB) noise (B and b), 1000 ± 50 parts per million (ppm) toluene (C and c), co-exposure to 100 ± 5 dB noise and 1000 ± 50 ppm toluene (D and d), and control group (A and a). Exposure routes: Inhalation exposure to toluene and hearing exposure to noise.

Fig. 6. Light micrograph of cortex lesions of frontal section stained with hematoxylin and eosin (H&E) at 10x (A, B, C, and D) and 40x (a, b, c, and d); Lesions were removed fourteen days subsequently the end of exposure to 100 ± 5 decibel (dB) noise (B and b), 1000 ± 50 parts per million (ppm) toluene (C and c), co-exposure to 100 ± 5 dB noise and 1000 ± 50 ppm toluene (D and d), and control group (A and a). Exposure routes: Inhalation exposure to toluene and hearing exposure to noise.

Fig. 7. Light micrograph of hippocampus lesions stained with hematoxylin and eosin (H&E) (40x); Lesions were removed fourteen days later the end of exposure to 100 ± 5 decibel (dB) noise (B), 1000 ± 50 parts per million (ppm) toluene (C), co-exposure to 100 ± 5 dB noise and 1000 ± 50 ppm toluene (D), and control group (A). The control group (A) observed without noticeable pathological changes. Some histopathological changes observed in the noise exposure group (B), in the toluene exposure group (C), and in the simultaneous exposure group (D) as marked in the figure. The signs marked in the photos presented pyramidal cells (dentate gyrus) (○) and dark cells (→). Exposure routes: Inhalation exposure to toluene and hearing exposure to noise