Title: Paternal Aggression in Early-Life Impairs the Spatial Memory and Passive Avoidance Learning in Adulthood of Male Rats: The Possible Role of DRD2

Running title: Ealy-life paternal aggression impairs learning memory

Authors: Solmaz Khalifeh^{1*}, Somayeh Tirbakhsh¹, Sareh Asadi², Ehsan Asadi², Ali Maleki², Fariba Khodaghali^{2*}, Mohammad-Reza Zarrindast^{3,4}, Mohammad Nasehi¹, Afshin Kheradmand⁵

1. Cognitive and Neuroscience Research Center (CNRC), Tehran Medical Sciences, Islamic Azad University, Tehran, Iran.
2. Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
3. Iranian National Center for Addiction Studies, Tehran University of Medical Sciences, Tehran, Iran.
4. Department of Pharmacology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.
5. Department of Pharmacology, School of Pharmacy-International Campus, Iran University of Medical Sciences, Tehran, Iran.

***Corresponding Authors:** 1. Solmaz Khalifeh, Cognitive and Neuroscience Research Center (CNRC), Tehran Medical Sciences, Islamic Azad University, Tehran, Iran. E-mail address: s.khalifeh@iautmu.ac.ir
2. Fariba Khodaghali, Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: Khodaghali@sbmu.ac.ir

To appear in: *Basic and Clinical Neuroscience*

Received date: 2021/06/10

Revised date: 2021/09/20

Accepted date: 2021/10/9

This is a “Just Accepted” manuscript, which has been examined by the peer-review process and has been accepted for publication. A “Just Accepted” manuscript is published online shortly after its acceptance, which is prior to technical editing and formatting and author proofing. *Basic and Clinical Neuroscience* provides “Just Accepted” as an optional and free service which allows authors to make their results available to the research community as soon as possible after acceptance. After a manuscript has been technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as a published article. Please note that technical editing may introduce minor changes to the manuscript text and/or graphics which may affect the content, and all legal disclaimers that apply to the journal pertain.

Please cite this article as:

Khalifeh, S., Tirbakhsh, S., Asadi, S., Asadi, E., Maleki, A., Khodagholi, F., et al. (In Press). Paternal Aggression in Early-Life Impairs the Spatial Memory and Passive Avoidance Learning in Adulthood of Male Rats: The Possible Role of DRD2. *Basic and Clinical Neuroscience*. Just Accepted publication Sep. 10, 2021. Doi:<http://dx.doi.org/10.32598/bcn.2021.3464.1>

DOI: <http://dx.doi.org/10.32598/bcn.2021.3464.1>

Abstract

Negative early-life experiences (e.g., having an aggressive father) can leave long-lasting impacts on the behavior. However, it is not clear if they influence learning and memory. In this study, we investigated the influences that the presence of an aggressive father had on the level of passive avoidance learning and spatial memory. We also studied the changes in the dopamine receptor D2 (DRD2) and peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α) gene expression in the hippocampus. Then, we evaluated if a DRD2 antagonist (Sulpiride, 0.125, 0.25, or 0.5 μ g/rat) could modulate these changes. We found that the subjects exposed to early-life stress made by aggressive fathers had impaired passive avoidance learning and spatial memory than those with normal fathers. Treatment with Sulpiride improved passive avoidance learning and spatial memory in rats with aggressive fathers. The rats with aggressive fathers also had higher expression of the DRD2 gene in their hippocampus than those with normal fathers, while the PGC-1 α gene expression was not different among groups. Treatment with Sulpiride (0.125, 0.25, or 0.5 μ g/rat) reduced the DRD2 gene expression in those with aggressive fathers to the normal level in those with normal fathers. These data suggest that living in a shared place with an aggressive father, even without any physical contact, can detrimentally affect passive avoidance learning and spatial memory which is accompanied by the increased expression of the DRD2 gene. Also, Sulpiride as a dopaminergic antagonist could reverse this process.

Keywords: Paternal aggression, Passive avoidance, Spatial memory, Sulpiride, Dopamine Receptor D2, PGC-1 α

1. Introduction

Learning and memory are two crucial physiological abilities by which animals can percept and restore information. These processes help animals recall the information in the moment of need and adjust the efficacy of their behaviors. Spatial memory has immense importance in the lives of animals as well as humans. For example, it helps animals to remember place of food and improve the outcome of foraging (Boyer & Walsh, 2010), associate positive and negative valences with various environments (Herry & Johansen, 2014; Kummer et al., 2011), and avoid potential predators.

Various internal and external factors can influence learning and memory capacity in animals. The early-life experience might be the most important modulator of the future brain's ability among the external factors. For example, experiencing a higher level of maternal care in the early-life period, as a positive environmental signal, improves spatial learning and memory (Liu, Diorio, Day, Francis, & Meaney, 2000). Not only the behavior of mothers, but also the behaviors that fathers show toward their offspring, like aggressive behavior, can have a deep impact on their behavioral development, like on their adulthood aggression (Frazier, Trainor, Cravens, Whitney, & Marler, 2006), depression, and anxiety (Khalifeh et al., 2020). However, the impact of being in a stressful condition with an aggressive father on the memory of offspring in adulthood is not very clear yet.

The literature suggests that dopamine receptor D2 (DRD2) plays an important role in spatial memory (Bezu et al., 2017; Raut, Jadhav, & Marathe, 2014). The activity of DRD2 can also modulate the performance of mice in the passive avoidance task (Ichihara, Nabeshima, & Kameyama, 1992). Peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α) is a key regulator of mitochondrial biogenesis (Valero, 2014), and also can be a modulator factor of learning and memory (Ashabi et al., 2012; Azimi, Gharakhanlou, Naghdi, Khodadadi, & Heysieattalab, 2018). Therefore, alterations in the level of DRD2 and PGC-1 α can modulate learning and memory. Early-life stress leaves long-lasting epigenetic marks, which switch off or on multiple genes (Silberman, Acosta, & Zorrilla Zubilete, 2016). This can be a ground for various behavioral impairments. The presence of an aggressive father can play as an early-life stressor. It raises whether the early-life presence of aggressive father can modulate later learning and memory by changing the expression of DRD2 and PGC-1 α genes. Moreover, Sulpiride has antidopaminergic properties and can antagonize DRD2 (Caley & Weber, 1995). It can modulate learning in rodents (Rinaldi, Mandillo, Oliverio, & Mele, 2007). Using these features, Sulpiride can help us to better understand the role of DRD2 in the scenario of having an aggressive father.

This study aimed to investigate whether the early-life presence of aggressive father can impact the passive avoidance learning and spatial memory of the male offspring in adulthood. We also questioned whether changes in DRD2 and PGC-1 α gene expression might have a role in this process. Then, we investigated the effects of Sulpiride treatment with three different doses in modulating the impact of aggressive fathers on learning and memory.

2. Materials and methods

2.1. Animals

Adult male Wistar rats were purchased from the Pasteur Institute (Tehran, Iran). They were kept in standard cages with temperature (22 ± 2 °C) and humidity-controlled environment in 12:12 h light/dark cycles (light on at 7:00 am) and were given unlimited access to water and standard rat chow *ad libitum*. All investigations and procedures were approved by the Research Ethics Committee of Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran (IR.SBMU.PHNS.REC.1397.055). During the entire process, highest effort was made to minimize the suffering of subjects.

2.2. Aggression training

Male albino Wistar rats were used as the resident rats based on the previously described method with some modifications (Hattori et al., 2015; Kinn Rød et al., 2014) (Fig. 1). Each male rat was housed with an adult female rat for two weeks. Before each session of testing the level of aggressiveness, the female rat was removed from the cage and an intruder male rat was placed into the cage. The behavior of the resident male rat was recorded for 5 minutes. Then, the intruder was removed, and the female was returned to the cage. The subjects that attack the intruder within the first 60 s of the session for three consecutive session and have higher attack latency in a shorter period of time are usually considered as aggressive (Miczek, 1979; Razzoli, Carboni, Andreoli, Ballottari, & Arban, 2011; Wood et al., 2015). Testing the level of aggressiveness was conducted for at least six sessions with four-day intervals. All the processes were performed during the light phase of the day. Resident rats with shorter attack latencies and more attack bites within a short period of time were considered as aggressive rats.

2.3. Mating

Both aggressive and non-aggressive male rats were used for mating. Each male rat was mated with a naïve non-aggressive female rat in a standard cage for 3 or 4 days and then removed.

2.4. Study design

After the birth of offspring, the biological fathers were returned to the home-cage for three weeks, but they were separated from females and offspring with a net, still having the olfactory, visual and auditory connections. Although physical contact was not possible, paternal olfactory, visual and auditory threats made a stressful atmosphere for offspring. Behavioral and molecular investigations were performed when the offspring were eight-week-old. In this study, male offspring of aggressive fathers were randomly divided into four treatment groups (n= 10/group) and the same division was done for male offspring of non-aggressive fathers (n= 10/group). In each aggressive and non-aggressive sets, the first group was just treated with saline (aggressive control group and non-aggressive control group). The next aggressive or non-aggressive groups received either 0.125, 0.25, or 0.5 µg/rat ICV Sulpiride treatment (Nasehi, Amin Yavari, & Zarrindast, 2013; Zarrindast, Nasehi, Pournaghshband, & Ghorbani Yekta, 2012). After that, the groups underwent behavioral experiments, namely passive avoidance task and Y-maze test. Then, the molecular studies were performed.

2.5. Drug preparation and surgery procedure

To prepare drug solution, Sulpiride (0.125, 0.25, or 0.5 µg/rat) was dissolved in 0.9 % physiological serum just before the injection. A combination of ketamine hydrochloride and xylazine were used for anesthetizing the offspring. Then, rats were prepared for surgery by stereotaxic device (Stoelting, USA). Guide cannula was inserted into the left ventricle (coordinates, 1.5mm lateral to the midline, – 0.8mm posterior to the bregma, and –4.6mm ventral to the skull surface). The injection (5 µl/rat) was done using Hamilton's syringe, and the drug or saline was injected slowly in 180 seconds.

2.6. Behavioral investigations

2.6.1. Passive avoidance task

The shuttle box was used to evaluate passive avoidance learning and memory (Brandewiede, Stork, & Schachner, 2014; Khakpour-Taleghani et al., 2008) with some modifications. The apparatus was made of polyvinyl chloride and consisted of two compartments separated with a removable gate; the dark compartment with a floor of electrically conductive rods and the light compartment. Electric shocks were delivered to the rods with an adjustable stimulator. For the first step, rats were habituated and trained in the apparatus. For the training, the observer put the rat in the light compartment and opened the gate ten seconds later. After the passage of four paws of rat from the gate, the observer closed the gate, and the rat was delivered an electrical shock (50 Hz, 1 mA intensity, 1.5 sec) in the dark compartment. If the rat did not enter the dark compartment after 120 seconds, it was considered a trained rat and ready for the main test.

Twenty-four hours after the training, passive avoidance memory retrieval test was carried out in all rats over which shock was not delivered (Pakdel & Rashidy-Pour, 2007). Similar to the training session, rats were put in the light compartment of the apparatus one by one. After opening the gate, the latency to enter the dark compartment for the first time and the time that rats spent in the dark compartment were recorded for ten minutes by the observer.

2.6.2. Y-maze test

Similar to the previously described methodology (Krauter, Guest, & Sarnyai, 2019) with some modifications, rats were tested in a Y-maze made of Plexiglass. It had three arms with a 120° angle between each arm. Rats were transferred to the test room 20 minutes before the beginning of trials. Then, they were placed in an arm of Y-maze which had a barrier. The barrier was removed 5 seconds after rats' entry to Y-maze, and all the movements of rats in different arms were recorded for eight minutes. Arm entry was considered as when all four paws of a rat passed the entrance of the arm. Based on the literature (Ansari Dezfouli, Zahmatkesh, Farahmandfar, & Khodagholi, 2019; Krauter et al., 2019), the successful consecutive entries as the index of behavioral test showing learning memory were calculated as “Successful consecutive entries = (Entries into three different arms consecutively / [number of total arm entries – 2]) × 100”.

2.7. Molecular analysis

After the behavioral tests, the rats were sacrificed, and the hippocampus was extracted for the molecular studies.

2.7.1. mRNA extraction and cDNA synthesis

We extracted total RNA from the hippocampus tissue using YTzol Pure RNA (Yektatajhez Azma, Tehran, Iran). The quality of RNA was investigated by ND-2000 spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA). Synthesis of cDNAs was done by Prime Script First Strand cDNA Synthesis Kit (Pars Tous, Tehran, Iran).

2.7.2. Real-time PCR

Real Time-PCR (RT-PCR) was performed using the Real Q-PCR Master Mix Kit (Ampliqon, Herlev, Denmark) on an ABI system (Step one). The samples were used to amplify PGC-1 α gene (forward primer, 5'-ATGAATGCAGCGGTCTTAGC-3'; reverse primer, 5'-AACAATGGCAGGGTTTGTTC-3'; annealing temperature 55 °C) and DRD2 gene (forward primer, 5'-CTGGTCGCCTCTTGTGC-3'; reverse primer, 5'-CGCATGACTCGTTCAGATCC-3'; annealing temperature 59 °C). For the control sample, distilled water was used. The β -actin gene (forward primer, 5'-TCTATCCTGGCCTCACTGTC-3'; reverse primer, 5'-AACGCAGCTCAGTAACAGTCC-3') was used as the housekeeping gene.

2.8. Statistical Analysis

Data were expressed as mean $\pm$ SEM and were analyzed using two-way ANOVA followed by Tukey-Test using Graph Pad Prism 5 software. The $p < 0.05$ was considered significant.

3. Results

3.1. Rats with aggressive fathers showed reduced passive avoidance learning and memory, and Sulpiride injection improved the outcome

Data indicated that for the time spent in the dark compartment, there were significant treatment effect ($p < 0.001$), group effect (aggressive vs. normal father) ($p < 0.001$), and treatment $\times$ group interaction ($p < 0.001$). The rats in the control group with aggressive fathers spent more time in the dark compartment of the passive avoidance task compared to the controls with normal fathers ($p < 0.001$) (Fig. 2a). Sulpiride injection in rats with aggressive fathers (all three doses: 0.125, 0.25, 0.5 μ g/rat) caused a significant decrease in the spent time in the dark part compared to the aggressive group (p s < 0.001). In addition, Sulpiride injection had no significant effect on the time that rats with normal fathers spent in the dark part compared to the control group. This measure was also different between rats with aggressive and normal fathers after injection of Sulpiride (0.125 μ g/rat) ($p < 0.001$). For the step-through latency, there were also significant treatment effect ($p < 0.001$), effect of aggressive father ($p < 0.001$), and treatment $\times$ aggressive father interaction ($p < 0.001$). Control rats with aggressive fathers had shorter step-through latency than the controls with normal fathers ($p < 0.001$) (Fig. 2b). Sulpiride injection in all three different doses (0.125, 0.25, 0.5 μ g/rat) caused a significant increase in the step-through latency in rats with the aggressive fathers compared to the group with aggressive fathers ($p < 0.001$), but did not have a significant

effect on rats with normal fathers compared to the controls with normal fathers. The step-through latency in rats with normal fathers treated with 0.25 µg/rat or 0.125 µg/rat Sulpiride was higher than those treated with aggressive fathers ($p < 0.001$).

3.2. Rats with aggressive fathers showed reduced spatial memory, but Sulpiride injection improved their memory

Y-maze test was used to evaluate spatial memory in rats of different groups. The results showed that for spontaneous alteration, there were significant treatment effect ($p < 0.001$), group effect (aggressive vs. normal father) ($p < 0.001$), and treatment $\times$ group interaction ($p < 0.001$). The control rats with aggressive fathers had less spontaneous alternation number than the controls with normal fathers ($p < 0.001$) (Fig. 3a). Sulpiride in all three different doses (0.125, 0.25, 0.5 µg/rat) increased the number of spontaneous alternations significantly in treated rats with aggressive fathers compared to the controls with aggressive fathers ($p < 0.001$). In contrast, Sulpiride injection in male rats with normal fathers made no significant change in spontaneous alternation compared to the controls with normal fathers.

We observed similar results for the total number of entries in Y-maze arms (Fig. 3b). For this parameter, there were significant treatment effect ($p < 0.001$), effect of aggressive father ($p < 0.001$), and treatment $\times$ aggressive father interaction ($p < 0.001$). The control rats with aggressive fathers had significantly less number of entries than those with normal fathers ($p < 0.001$), but Sulpiride injection in all three doses (0.125, 0.25, and 0.5 µg/rat) caused a significant increase ($p < 0.001$). Also, Sulpiride injection in male rats with normal fathers made no significant change in the total number of entries in treated rats with normal fathers compared to the controls with normal fathers.

3.3. Rats with aggressive fathers had a higher level of DRD2 gene expression in the hippocampus, and Sulpiride injection reversed this effect

Results from RT-PCR showed that there were significant treatment effect ($p < 0.001$), effect of aggressive father ($p < 0.001$), and treatment $\times$ aggressive father interaction ($p < 0.001$) on the level of DRD2 gene expression in the hippocampus. The DRD2 gene expression of the controls with aggressive fathers was significantly higher compared to the controls with normal fathers ($p < 0.001$) (Fig. 4). Sulpiride injection (0.125, 0.25, 0.5 µg/rat) in treated rats with aggressive fathers significantly decreased the level of DRD2 mRNAs compared to the controls with aggressive fathers ($p < 0.001$). Sulpiride injection in rats with normal fathers made no significant change in DRD2 gene expression level compared to the controls with normal fathers.

3.4. Rats with aggressive fathers did not have different PGC-1 α gene expression in the hippocampus, and Sulpiride injection also did not cause any change

We observed no significant treatment effect, effect of aggressive father, or treatment $\times$ aggressive father interaction. PGC-1 α gene expression was not different between controls with aggressive fathers and controls with

normal fathers (Fig. 5). Also, Sulpiride treatment with either dosage did not change the level of PGC-1 α gene expression compared to the control rats with either aggressive or normal fathers.

4. Discussion

In this study, we aimed to investigate the consequences of paternal aggressiveness on their offspring's learning and memory. We found that even though there was no physical contact with an aggressive father, offspring exposure to this environment made a challenging atmosphere in which baby rats' passive avoidance learning and spatial memory were impaired. Moreover, the rats being in this environment had significantly higher DRD2 gene expression in the hippocampus compared to the normal rats, while the PGC-1 α gene expression did not significantly differ. The rats with aggressive fathers that were treated with different dosages of Sulpiride (0.125, 0.25, or 0.5 μ g/rat) showed significantly improved passive avoidance learning and spatial memory compared to the controls. The treatment also normalized the expression of the DRD2 gene in offspring with aggressive fathers to the level of control group.

Ample human evidence showed that children's exposure to domestic violence leads to long-term psychiatric and behavioral detriments (Cicchetti, 2016; Marshall, Feinberg, & Daly, 2019; Vachon, Krueger, Rogosch, & Cicchetti, 2015). Indeed, domestic violence and exposure to this environment are believed to be a precursor to a wide range of psychological disorders for children (Margolin & Vickerman, 2007). Majali et al., found that children as witnesses or victims of parental aggression can show psychological and emotional disturbances in their adulthoods (Al Majali & Alsrehan, 2019). Socio-psychological disability, stress, anxiety, and less self-esteem have been reported in these children (Al Majali & Alsrehan, 2019; Øverlien, 2010; Rehan, Antfolk, Johansson, & Santtila, 2019). Chronic stress is one the most common consequences of parental aggressiveness which detrimentally affects children's physical and mental health (Moffitt, 2013). Chronic stress as a result of domestic violence can lead to psychological alterations in some regions of brain such as cerebral cortex, limbic system, corpus callosum, cerebellum, and hypothalamus (Tsavoussis, Stawicki, Stoicea, & Papadimos, 2014). Smith and Pollak. also found that early life stress can have a permanent and pervasive impact on the hippocampus, amygdala, prefrontal network, and dopaminergic circuits (Smith & Pollak, 2020). Negative emotions activate amygdala-dependent memory formation and change the associative memory performance (Okada et al., 2011). In a study by Nguyen., et al, it has been revealed that the environment in which children live strongly affects their emotions, behaviors, and memories. Negative emotions and unpleasant behaviors in children resulted from aggressive parents in early postnatal life lead to impaired hippocampal-dependent memory, aggression, anxiety, and fear-memory formation in adulthood (Nguyen, Bagot, Diorio, Wong, & Meaney, 2015). This underlines the long-lasting impact of early-life paternal behavior on offspring.

In this study, control rats experiencing a traumatic and anxious environment with aggressive fathers had shorter step-through latencies and spent more time in the dark compartment of the passive avoidance test. Our results showed that the passive avoidance learning and memory of these rats were reduced compared to rats with normal

situation. In addition, in the Y-maze test, control rats being under pressure with an aggressive father had less successful spontaneous alternations and fewer entries than the control groups living in a healthy environment, which means they have reduced spatial memory compared to rats with normal conditions. Therefore, these findings support the negative effects of early-life paternal aggression on passive avoidance learning and spatial memory of offspring in adulthood. This might be due to the early-life stress induced by the paternal aggression and the related memory impairments. For example, early-life stress induced by other sources like maternal separation can also impair learning and memory (Maghami et al., 2018), which supports the stress as the possible bridge between paternal aggression and impaired learning and memory.

DRD2, as a member of the dopaminergic receptors family, affects the spatial and long-term memory in rats (Rocchetti et al., 2015). Different studies have shown different associations between this receptor and negative emotions such as depression (Chiesa et al., 2014), anxiety (Wang et al., 2014), and fear (Huertas et al., 2010). The role of this gene in learning and memory have also been demonstrated (Rocchetti et al., 2015; Zhu et al., 2010). It is also involved in parental care (Curry et al., 2013; Zhu et al., 2010). In 2013, Yi Zhang et al. found that maternal separation- which also is an early-life stressor - results in changes in the DRD2 gene expression in nucleus accumbens, which leads to behavioral problems and spatial memory impairment in adulthood (Zhang et al., 2013). We found that rats that experienced tension in their early life with aggressive fathers had a higher level of DRD2 gene expression compared to the rats with normal situation. This result is similar to other studies which demonstrated an association between memory impairment and increased expression of DRD2 gene (Noohi et al., 2014; Persson, Rieckmann, Kalpouzos, Fischer, & Bäckman, 2015). We also found that i.c.v Sulpiride injection in either three different doses can improve the passive avoidance and spatial memory in rats with aggressive fathers.

Moreover, stress can increase the DRD2 gene expression in memory-related brain regions like prefrontal cortex (Tomas-Roig & Havemann-Reinecke, 2019). In specific, perinatal stress can increase DRD2 gene expression in the hippocampus of rats (Berger, Barros, Sarchi, Tarazi, & Antonelli, 2002). This is similar to our findings that paternal aggression in early life (as a stressor) increases DRD2 expression in the hippocampus. In terms of the roles that this receptor plays in the memory, literature suggests that when the level of hippocampal DRD2 is in the physiologic range, its activation may improve and its deactivation may impair learning and memory (Wilkerson & Levin, 1999). However, hyperactivation of DRD2 in the hippocampus during adolescent impairs spine development, neural circuits, and spatial memory. Interestingly, DRD2 blockers prevented negative impacts of DRD2 hyperactivation on spine development and spatial memory (Jia, Zhao, Hu, Lindberg, & Li, 2013). These evidences might explain our findings. The early-life experience of aggressive father may act as a stressor and, therefore, increase the DRD2 gene expression level in the hippocampus. This increased level may cause hyperactivation of DRD2 receptors and impairs spatial memory by interfering with spine development of the neurons. As we observed, the memory impairment was reversed by using Sulpiride, as a DRD2 blocker.

On the other side, Sulpiride, as an antagonist of DRD2 in the central nervous system, is used for the treatment of psychiatric diseases (Caley & Weber, 1995; Ciszowski, Szpak, & Wilimowska, 2010). In line with our results, Floresco et al. found that Sulpiride can be a promising candidate for the treatment of various psychological disorders, enhancement of memory and improvement of attention, working memory, and behavioral flexibility (Floresco & Jentsch, 2011). Similarly, Passetti et al., showed that medial prefrontal cortex impairment is relevant to neuropsychological dysfunction. on the other hand, modulating the dopaminergic pathway in basal ganglia can alleviate any neuropsychological deficit (Passetti, Levita, & Robbins, 2003). In 2005, Mehta et al. reported that Sulpiride can improve working memory and facilitate the responses to the visual learning tasks (Mehta, Hinton, Montgomery, Bantick, & Grasby, 2005). The half-life of Sulpiride is 24 hours in the case of systemic injection, but we followed the literature and used i.c.v injection to increase its lifetime and provide better availability of the drug in the brain (Mizuchi, Kitagawa, & Miyachi, 1983).

PGC-1 α is a key regulator of mitochondrial biogenesis (Valero, 2014), and also can be a modulator factor of learning and memory (Ashabi et al., 2012; Azimi, Gharakhanlou, Naghdi, Khodadadi, & Heysieattalab, 2018). It is shown that acute stress in the adulthood can change learning and memory by modulating the level of PGC-1 α (Sardari, Rezayof, & Khodagholi, 2015). However, our observation showed that the PGC-1 α gene expression in the rats with aggressive fathers was not significantly different from those with normal fathers. It suggests that, in contrast to acute stress in adulthood, the early-life stress of having an aggression father do not change the PGC-1 α gene expression in the hippocampus. The Sulpiride treatment did not change this expression. Moreover, the injection of Sulpiride with different doses in rats with normal fathers had no significant effect on their passive avoidance and spatial memory as well as their DRD2 and PGC-1 α gene expression profile.

These results of our study showed that living in a shared place with an aggressive father, even without any physical contact, can detrimentally affect passive avoidance learning and spatial memory which is accompanied by the increased expression of the DRD2 gene. However, Sulpiride as a dopaminergic antagonist could reverse this process.

Ethical Considerations

Compliance with ethical guidelines

All investigations and procedures were approved by the Research Ethics Committee of Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran (IR.SBMU.PHNS.REC.1397.055).

Funding

This work was supported by Shahid Beheshti University of Medical Sciences Research h-index grant (Grant No. 14945-7-3).

Conflict of interest

The authors had no conflict of interest to declare.

Acknowledgments

This work was supported by Shahid Beheshti University of Medical Sciences Research h-index grant (Grant No. 14945-7-3).

Figure legends

Fig 1. The timeline that the procedures were conducted.

Fig 2. Passive avoidance learning and memory using passive avoidance task. **A** The time spent in the dark compartment in different groups of rats. **B** The time of first entrance to the dark compartment of the task in different groups of rats. Each bar shows the mean $\pm$ SEM. (n = 10/group). ###p < 0.001 vs. controls with normal fathers, ***p < 0.001 vs. controls with aggressive fathers, &&&p < 0.001.

Fig 3. Spatial memory using Y-maze. **A** The number of spontaneous alternations in the Y-maze test in different groups of rats. **B** The total number of movements in arms in the Y-maze test in different groups of rats. Each bar shows the mean $\pm$ SEM. (n = 10/group). ###p < 0.001 vs. controls with normal fathers, ***p < 0.001 vs. controls with aggressive fathers.

Fig 4. The level of DRD2 gene expression in the hippocampus of different groups of rats. Each point shows the mean $\pm$ SEM. (n = 4/group). ###p < 0.001 vs. controls with normal fathers, ***p < 0.001 vs. controls with aggressive fathers.

Fig 5. The level of PGC-1 α gene expression in the hippocampus of different groups of rats. Each point shows the mean $\pm$ SEM. (n = 4/group).

A.

B.

A.

B.

References

- Al Majali, S., & Alsrehan, H. (2019). The impact of family violence on the social and psychological development of the child. *Utopía y Praxis Latinoamericana*, 24(5), 199-207.
- Ansari Dezfouli, M., Zahmatkesh, M., Farahmandfar, M., & Khodagholi, F. (2019). Melatonin protective effect against amyloid beta-induced neurotoxicity mediated by mitochondrial biogenesis; involvement of hippocampal Sirtuin-1 signaling pathway. *Physiol Behav*, 204, 65-75. doi:10.1016/j.physbeh.2019.02.016
- Ashabi, G., Ramin, M., Azizi, P., Taslimi, Z., Alamdary, S. Z., Haghparast, A., . . . Khodagholi, F. (2012). ERK and p38 inhibitors attenuate memory deficits and increase CREB phosphorylation and PGC-1alpha levels in Abeta-injected rats. *Behav Brain Res*, 232(1), 165-173. doi:10.1016/j.bbr.2012.04.006
- Azimi, M., Gharakhanlou, R., Naghdi, N., Khodadadi, D., & Heysieattalab, S. (2018). Moderate treadmill exercise ameliorates amyloid-beta-induced learning and memory impairment, possibly via increasing AMPK activity and up-regulation of the PGC-1alpha/FNDC5/BDNF pathway. *Peptides*, 102, 78-88. doi:10.1016/j.peptides.2017.12.027
- Berger, M. A., Barros, V. G., Sarchi, M. I., Tarazi, F. I., & Antonelli, M. C. (2002). Long-term effects of prenatal stress on dopamine and glutamate receptors in adult rat brain. *Neurochem Res*, 27(11), 1525-1533. doi:10.1023/a:1021656607278
- Bezu, M., Malikovic, J., Kristofova, M., Engidawork, E., Hoger, H., Lubec, G., & Korz, V. (2017). Spatial Working Memory in Male Rats: Pre-Experience and Task Dependent Roles of Dopamine D1- and D2-Like Receptors. *Front Behav Neurosci*, 11, 196. doi:10.3389/fnbeh.2017.00196
- Boyer, D., & Walsh, P. D. (2010). Modelling the mobility of living organisms in heterogeneous landscapes: does memory improve foraging success? *Philos Trans A Math Phys Eng Sci*, 368(1933), 5645-5659. doi:10.1098/rsta.2010.0275
- Brandewiede, J., Stork, O., & Schachner, M. (2014). NCAM deficiency in the mouse forebrain impairs innate and learned avoidance behaviours. *Genes, Brain and Behavior*, 13(5), 468-477.
- Caley, C. F., & Weber, S. S. (1995). Sulpiride: an antipsychotic with selective dopaminergic antagonist properties. *Ann Pharmacother*, 29(2), 152-160. doi:10.1177/106002809502900210
- Chiesa, A., Lia, L., Alberti, S., Lee, S.-J., Han, C., Patkar, A. A., . . . Serretti, A. (2014). Lack of influence of rs4680 (COMT) and rs6276 (DRD2) on diagnosis and clinical outcomes in patients with major depression. *International Journal of Psychiatry in Clinical Practice*, 18(2), 97-102. doi:10.3109/13651501.2014.894073
- Cicchetti, D. (2016). Socioemotional, personality, and biological development: Illustrations from a multilevel developmental psychopathology perspective on child maltreatment. *Annual review of psychology*, 67, 187-211.

- Ciszowski, K., Szpak, D., & Wilimowska, J. (2010). [Toxicity of sulpiride]. *Przegląd Lekarski*, 67(8), 606-609. Retrieved from <http://europepmc.org/abstract/MED/21387787>
- Curry, T., Egeto, P., Wang, H., Podnos, A., Wasserman, D., & Yeomans, J. (2013). Dopamine receptor D2 deficiency reduces mouse pup ultrasonic vocalizations and maternal responsiveness. *Genes, Brain & Behavior*, 12(4), 397-404. doi:10.1111/gbb.12037
- Floresco, S. B., & Jentsch, J. D. (2011). Pharmacological enhancement of memory and executive functioning in laboratory animals. *Neuropsychopharmacology*, 36(1), 227-250.
- Frazier, C. R., Trainor, B. C., Cravens, C. J., Whitney, T. K., & Marler, C. A. (2006). Paternal behavior influences development of aggression and vasopressin expression in male California mouse offspring. *Horm Behav*, 50(5), 699-707. doi:10.1016/j.yhbeh.2006.06.035
- Hattori, T., Kanno, K., Nagasawa, M., Nishimori, K., Mogi, K., & Kikusui, T. (2015). Impairment of interstrain social recognition during territorial aggressive behavior in oxytocin receptor-null mice. *Neuroscience Research*, 90, 90-94. doi:<https://doi.org/10.1016/j.neures.2014.05.003>
- Herry, C., & Johansen, J. P. (2014). Encoding of fear learning and memory in distributed neuronal circuits. *Nat Neurosci*, 17(12), 1644-1654. doi:10.1038/nn.3869
- Huertas, E., Ponce, G., Koeneke, M. A., Poch, C., España-Serrano, L., Palomo, T., . . . Hoenicka, J. (2010). The D2 dopamine receptor gene variant C957T affects human fear conditioning and aversive priming. *Genes Brain Behav*, 9(1), 103-109. doi:10.1111/j.1601-183X.2009.00543.x
- Ichihara, K., Nabeshima, T., & Kameyama, T. (1992). Effects of dopamine receptor agonists on passive avoidance learning in mice: interaction of dopamine D1 and D2 receptors. *Eur J Pharmacol*, 213(2), 243-249. doi:10.1016/0014-2999(92)90688-z
- Jia, J. M., Zhao, J., Hu, Z., Lindberg, D., & Li, Z. (2013). Age-dependent regulation of synaptic connections by dopamine D2 receptors. *Nat Neurosci*, 16(11), 1627-1636. doi:10.1038/nn.3542
- Khakpour-Taleghani, B., Lashgari, R., Aavani, T., Haghparast, A., Naderi, N., & Motamedi, F. (2008). The locus coeruleus involves in consolidation and memory retrieval, but not in acquisition of inhibitory avoidance learning task. *Behav Brain Res*, 189(2), 257-262. doi:10.1016/j.bbr.2008.01.004
- Khalifeh, S., Khodagholi, F., Zarrindast, M. R., Alizadeh, R., Asadi, S., Mohammadi Kamsorkh, H., . . . Sadat-Shirazi, M. S. (2020). Altered D2 receptor and transcription factor EB expression in offspring of aggressive male rats, along with having depressive and anxiety-like behaviors. *Int J Neurosci*, 1-11. doi:10.1080/00207454.2020.1758086
- Kinn Rød, A. M., Murison, R., Mrdalj, J., Milde, A. M., Jellestad, F. K., Øvernes, L. A., & Grønli, J. (2014). Effects of social defeat on sleep and behaviour: Importance of the confrontational behaviour. *Physiology & Behavior*, 127, 54-63. doi:<https://doi.org/10.1016/j.physbeh.2014.01.010>

- Kraeuter, A. K., Guest, P. C., & Sarnyai, Z. (2019). The Y-Maze for Assessment of Spatial Working and Reference Memory in Mice. *Methods Mol Biol*, 1916, 105-111. doi:10.1007/978-1-4939-8994-2_10
- Kummer, K., Klement, S., Eggart, V., Mayr, M. J., Saria, A., & Zernig, G. (2011). Conditioned place preference for social interaction in rats: contribution of sensory components. *Front Behav Neurosci*, 5, 80. doi:10.3389/fnbeh.2011.00080
- Liu, D., Diorio, J., Day, J. C., Francis, D. D., & Meaney, M. J. (2000). Maternal care, hippocampal synaptogenesis and cognitive development in rats. *Nat Neurosci*, 3(8), 799-806. doi:10.1038/77702
- Maghami, S., Zardooz, H., Khodagholi, F., Binayi, F., Ranjbar Saber, R., Hedayati, M., . . . Ansari, M. A. (2018). Maternal separation blunted spatial memory formation independent of peripheral and hippocampal insulin content in young adult male rats. *PLoS One*, 13(10), e0204731. doi:10.1371/journal.pone.0204731
- Margolin, G., & Vickerman, K. A. (2007). Post-traumatic Stress in Children and Adolescents Exposed to Family Violence: I. Overview and Issues. *Prof Psychol Res Pr*, 38(6), 613-619. doi:10.1037/0735-7028.38.6.613
- Marshall, A. D., Feinberg, M. E., & Daly, K. A. (2019). Children's emotional and behavioral reactions to interparental aggression: The role of exposure to within-incident, cross-dyad aggression spillover. *Journal of family psychology*, 33(5), 617.
- Mehta, M. A., Hinton, E. C., Montgomery, A. J., Bantick, R. A., & Grasby, P. M. (2005). Sulpiride and mnemonic function: effects of a dopamine D2 receptor antagonist on working memory, emotional memory and long-term memory in healthy volunteers. *J Psychopharmacol*, 19(1), 29-38. doi:10.1177/0269881105048889
- Miczek, K. A. (1979). A new test for aggression in rats without aversive stimulation: differential effects of d-amphetamine and cocaine. *Psychopharmacology (Berl)*, 60(3), 253-259. doi:10.1007/BF00426664
- Mizuchi, A., Kitagawa, N., & Miyachi, Y. (1983). Regional distribution of sultopride and sulpiride in rat brain measured by radioimmunoassay. *Psychopharmacology (Berl)*, 81(3), 195-198. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/6417707>
- Moffitt, T. E. (2013). Childhood exposure to violence and lifelong health: clinical intervention science and stress-biology research join forces. *Dev Psychopathol*, 25(4 Pt 2), 1619-1634. doi:10.1017/s0954579413000801
- Nasehi, M., Amin Yavari, S., & Zarrindast, M. R. (2013). Synergistic effects between CA1 mu opioid and dopamine D1-like receptors in impaired passive avoidance performance induced by hepatic encephalopathy in mice. *Psychopharmacology*, 227(3), 553-566. doi:10.1007/s00213-013-2987-y
- Nguyen, H.-B., Bagot, R. C., Diorio, J., Wong, T. P., & Meaney, M. J. (2015). Maternal care differentially affects neuronal excitability and synaptic plasticity in the dorsal and ventral hippocampus. *Neuropsychopharmacology*, 40(7), 1590-1599.

- Noohi, F., Boyden, N. B., Kwak, Y., Humfleet, J., Burke, D. T., Müller, M. L., . . . Seidler, R. D. (2014). Association of COMT val158met and DRD2 G>T genetic polymorphisms with individual differences in motor learning and performance in female young adults. *J Neurophysiol*, 111(3), 628-640. doi:10.1152/jn.00457.2013
- Okada, G., Okamoto, Y., Kunisato, Y., Aoyama, S., Nishiyama, Y., Yoshimura, S., . . . Yamawaki, S. (2011). The effect of negative and positive emotionality on associative memory: an fMRI study. *PLoS One*, 6(9), e24862.
- Øverlien, C. (2010). Children exposed to domestic violence: Conclusions from the literature and challenges ahead. *Journal of social work*, 10(1), 80-97.
- Pakdel, R., & Rashidy-Pour, A. (2007). Microinjections of the dopamine D2 receptor antagonist sulpiride into the medial prefrontal cortex attenuate glucocorticoid-induced impairment of long-term memory retrieval in rats. *Neurobiology of learning and memory*, 87(3), 385-390.
- Passetti, F., Levita, L., & Robbins, T. W. (2003). Sulpiride alleviates the attentional impairments of rats with medial prefrontal cortex lesions. *Behav Brain Res*, 138(1), 59-69. doi:10.1016/s0166-4328(02)00229-2
- Persson, J., Rieckmann, A., Kalpouzos, G., Fischer, H., & Bäckman, L. (2015). Influences of a DRD2 polymorphism on updating of long-term memory representations and caudate BOLD activity: magnification in aging. *Human brain mapping*, 36(4), 1325-1334. doi:10.1002/hbm.22704
- Raut, S. B., Jadhav, K. S., & Marathe, P. A. (2014). Role of dopamine--D2 receptor in spatial memory retention and retrieval determined using Hebb-Williams complex maze. *Indian J Physiol Pharmacol*, 58(3), 192-196. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/25906600>
- Razzoli, M., Carboni, L., Andreoli, M., Ballottari, A., & Arban, R. (2011). Different susceptibility to social defeat stress of BalbC and C57BL6/J mice. *Behav Brain Res*, 216(1), 100-108. doi:10.1016/j.bbr.2010.07.014
- Rehan, W., Antfolk, J., Johansson, A., & Santtila, P. (2019). Do Single Experiences of Childhood Abuse Increase Psychopathology Symptoms in Adulthood? *J Interpers Violence*, 34(5), 1021-1038. doi:10.1177/0886260516647004
- Rinaldi, A., Mandillo, S., Oliverio, A., & Mele, A. (2007). D1 and D2 receptor antagonist injections in the prefrontal cortex selectively impair spatial learning in mice. *Neuropsychopharmacology*, 32(2), 309-319. doi:10.1038/sj.npp.1301176
- Rocchetti, J., Isingrini, E., Dal Bo, G., Sagheby, S., Menegaux, A., Tronche, F., . . . Giros, B. (2015). Presynaptic D2 Dopamine Receptors Control Long-Term Depression Expression and Memory Processes in the Temporal Hippocampus. *Biological Psychiatry*, 77(6), 513-525. doi:<https://doi.org/10.1016/j.biopsych.2014.03.013>
- Sardari, M., Rezayof, A., & Khodagholi, F. (2015). Hippocampal signaling pathways are involved in stress-induced impairment of memory formation in rats. *Brain Res*, 1625, 54-63. doi:10.1016/j.brainres.2015.08.015

- Silberman, D. M., Acosta, G. B., & Zorrilla Zubilete, M. A. (2016). Long-term effects of early life stress exposure: Role of epigenetic mechanisms. *Pharmacol Res*, 109, 64-73. doi:10.1016/j.phrs.2015.12.033
- Smith, K. E., & Pollak, S. D. (2020). Early life stress and development: potential mechanisms for adverse outcomes. *Journal of Neurodevelopmental Disorders*, 12(1), 1-15.
- Tomas-Roig, J., & Havemann-Reinecke, U. (2019). Gene expression signature in brain regions exposed to long-term psychosocial stress following acute challenge with cannabinoid drugs. *Psychoneuroendocrinology*, 102, 1-8. doi:10.1016/j.psyneuen.2018.11.023
- Tsavoussis, A., Stawicki, S. P., Stoicea, N., & Papadimos, T. J. (2014). Child-witnessed domestic violence and its adverse effects on brain development: a call for societal self-examination and awareness. *Front Public Health*, 2, 178. doi:10.3389/fpubh.2014.00178
- Vachon, D. D., Krueger, R. F., Rogosch, F. A., & Cicchetti, D. (2015). Assessment of the harmful psychiatric and behavioral effects of different forms of child maltreatment. *JAMA psychiatry*, 72(11), 1135-1142.
- Valero, T. (2014). Mitochondrial biogenesis: pharmacological approaches. *Curr Pharm Des*, 20(35), 5507-5509. doi:10.2174/138161282035140911142118
- Wang, Y. S., Lee, S. Y., Chen, S. L., Chang, Y. H., Wang, T. Y., Lin, S. H., . . . Lu, R. B. (2014). Role of DRD2 and ALDH2 genes in bipolar II disorder with and without comorbid anxiety disorder. *European Psychiatry*, 29(3), 142-148. doi:<https://doi.org/10.1016/j.eurpsy.2013.05.001>
- Wilkerson, A., & Levin, E. D. (1999). Ventral hippocampal dopamine D1 and D2 systems and spatial working memory in rats. *Neuroscience*, 89(3), 743-749. doi:10.1016/s0306-4522(98)00346-7
- Wood, S. K., Wood, C. S., Lombard, C. M., Lee, C. S., Zhang, X. Y., Finnell, J. E., & Valentino, R. J. (2015). Inflammatory Factors Mediate Vulnerability to a Social Stress-Induced Depressive-like Phenotype in Passive Coping Rats. *Biol Psychiatry*, 78(1), 38-48. doi:10.1016/j.biopsych.2014.10.026
- Zarrindast, M. R., Nasehi, M., Pournaghshband, M., & Ghorbani Yekta, B. (2012). Dopaminergic system in CA1 modulates MK-801 induced anxiolytic-like responses. *Pharmacology Biochemistry and Behavior*, 103(1), 102-110. doi:<https://doi.org/10.1016/j.pbb.2012.07.016>
- Zhang, Y., Zhu, X., Bai, M., Zhang, L., Xue, L., & Yi, J. (2013). Maternal deprivation enhances behavioral vulnerability to stress associated with miR-504 expression in nucleus accumbens of rats. *PLoS One*, 8(7), e69934. doi:10.1371/journal.pone.0069934
- Zhu, X., Li, T., Peng, S., Ma, X., Chen, X., & Zhang, X. (2010). Maternal deprivation-caused behavioral abnormalities in adult rats relate to a non-methylation-regulated D2 receptor levels in the nucleus accumbens. *Behavioural Brain Research*, 209(2), 281-288. doi:<https://doi.org/10.1016/j.bbr.2010.02.005>

Accepted Manuscript (Uncorrected Proof)